



Antarctic Clouds simulated by Polar WRF and AMPS


Keith M. Hines, David H. Bromwich
and Sheng-Hung Wang

*Polar Meteorology Group
Byrd Polar and Climate Research Center
The Ohio State University
Columbus, OH*



U.S. DEPARTMENT OF
ENERGY

Office of
Science






ARM West Antarctic Radiation Experiment (AWARE)

OBSERVATIONS: 23 NOVEMBER 2015 - 5 JANUARY 2017

**Observations at
West Antarctic
Divide (WAIS) and
McMurdo**



**West Antarctic
warming during
January 2016**



AWARE aims to gain insight into the factors behind recent climate change in West Antarctica by quantifying the role of changing air masses on the surface energy balance. The field campaign use some of the most advanced atmospheric research instrumentation to conduct cloud, radiative, and aerosol observations.

Important for AWARE and WRF

**Does our knowledge of Arctic clouds
carry over to high southern latitudes?**


**Polar cloud observations (and modeling
studies) much more extensive in Arctic than
Antarctic**


Antarctica is colder, drier and more pristine

**AWARE observations suggest clouds less
extensive than over the Arctic basin, however
vertical motions are larger (topography?)**


Cloud water at very low temperature

Vertical cloud distribution is different





The Antarctic Mesoscale Prediction System (AMPS)



- **Adapted numerical weather prediction system for Antarctica**
 - Polar WRF (Weather Research and Forecasting Model)
 - Variable resolution now to 0.9 km
- **Priority Mission: U.S. Antarctic Program (USAP) Weather Support (clouds important for aircraft!)**
- **Collaborators: NCAR and OSU BPCRC**
- **Powers et al. (2012) A decade of Antarctic science through AMPS. BAMS, 93, 1699-1712.**
- **<http://www.mmm.ucar.edu/rt/amps>**



NCAR






NCAR

AMPS GRIDS



**Use AMPS
grid 2 for
WAIS
evaluation**



**Use December
2015 and
January 2016
AMPS forecasts
and WAIS
observations**








Test West Antarctic Summer Results for AMPS with WAIS Observations

Surface Energy Balance: Excess shortwave and deficit in longwave → Cloud deficit?


T-test



AMPS shows biases suggesting a better Antarctic cloud simulation is needed



Test Microphysics Schemes vs. WAIS Observations



**PWRF 3.9.1 on AMPS Grid 2 (10 km) with ERA-I
I.C. + B.C. (AMPS uses GFS)**

WRF Single-Moment 5-Class (same as AMPS)

Morrison 2-Moment (slight polar modifications)


Thompson-Eidhammer Aerosol AWARE

**Morrison-Milbrandt P3 (avoids arbitrary cloud
and precipitation categorization)**

**Note: New PWRF 3.9.1 runs have 24 hr spin-up
AMPS has 12-hr spin-up**




Near Surface Fields at WAIS 8 – 15 January 2016



PWRF 3.9.1: SW and LW biases remain, but are reduced in magnitude. Temperature and humidity biases are largely removed. Can use PWRF 3.9.1 to explore Antarctic cloud biases (AMPS linked).



Near Surface Fields at WAIS 8 – 15 January 2016







**Run with more advanced microphysics schemes: Warm bias in 2-m T?
Schemes increase LW and reduce SW radiation – positive result!**



Average PWRF 3.9.1 Forecasts

Cloud Forcing at WAIS 8 – 15 January 2016




WSM5C has smallest LWP and slow spin-up of longwave cloud forcing.
M-M P3 scheme has largest LWP and stronger SWCF.
Microphysics schemes impact cloud radiative effects for Antarctica!



Cloud Condensate Path at WAIS 8 – 15 January 2016

AMPS and PWRF 3.9.1 with WRF Single-Moment 5-Class (WSM5C)



AMPS and PWRF 3.9.1 with MSM5C microphysics show much more ice water path than liquid water path.


Simulated LWP is much less than values measured by the microwave radiometer (MWR).

Liquid has greater impact on radiation than ice.

Antarctic observations show liquid cloud at very low temperature.



Cloud Condensate Path at WAIS 8 – 15 January 2016



More advanced microphysics schemes increase ice water path and greatly increase liquid water path.

Morrison-Milbrandt P3 scheme shows a spike in liquid water on 11 January.

Day-to-day match of simulated and observed LWP is poor.

Simulating cloud water on cold days needs research.



Summary of AMPS and PWRP Findings with the AWARE Project

Liquid water deficit in AMPS clouds

Cloud radiative effect of AMPS clouds is too small

More advanced microphysics schemes increase the simulated liquid water and increase the cloud radiative effect

Which microphysics scheme is best? – not certain yet

Need to work on simulating cloud water at colder temperatures.

Clouds are critical for improving AMPS