

WRF Physics Options

Jimmy Dudhia

WRF Physics

- ◆ Turbulence/Diffusion (diff_opt, km_opt)
- ◆ Radiation
 - Longwave (ra_lw_physics)
 - Shortwave (ra_sw_physics)
- ◆ Surface
 - Surface layer (sf_sfclay_physics)
 - Land/water surface (sf_surface_physics)
- ◆ PBL (bl_physics)
- ◆ Cumulus parameterization (cu_physics)
- ◆ Microphysics (mp_physics)

Turbulence/Diffusion

Sub-grid eddy mixing effects on
all fields

diff_opt=1

◆ 2nd order diffusion on model levels

- Constant vertical coefficient (kvdif)
- Or Use with PBL

◆ km_opt

- 1: constant (khdif and kvdif used)
- 2: 1.5-order TKE prediction (not recommended with diff_opt=1)
- 3: Smagorinsky (deformation/stability based K) (not recommended with diff_opt=1)
- 4: 2D Smagorinsky (deformation based on horizontal wind for horizontal diffusion only)

diff_opt=2

- ◆ 2nd order horizontal diffusion
- ◆ Allows for terrain-following coordinate
- ◆ km_opt
 - 1: constant (khdif and kvdif used)
 - 2: 1.5-order TKE prediction
 - 3: Smagorinsky (deformation/stability based K)
 - 4: 2D Smagorinsky (deformation based on horizontal wind for horizontal diffusion only)

diff_opt=2 (continued)

- ◆ `mix_full_fields=.true.:` vertical diffusion acts on full (not perturbation) fields (recommended)
- ◆ Idealized constant surface fluxes can be added in `diff_opt=2` using namelist (dynamics section)
 - `tke_drag_coefficient` (C_D)
 - `tke_heat_flux` ($=H/\rho c_p$)

Diffusion Option Choice

- ◆ Real-data case with PBL physics on
 - Best is diff_opt=1, km_opt=4
 - This complements vertical diffusion done by PBL scheme
- ◆ Idealized large-eddy resolving cases
 - km_opt=2 (tke scheme) is designed for hi-res eddy-resolving modeling
- ◆ Cloud-resolving modeling (smooth or no topography)
 - diff_opt=1; km_opt=2,3
- ◆ Complex topography
 - diff_opt=2 is more accurate for sloped coordinate surfaces, and prevents diffusion up/down valley sides
- ◆ Note: WRF can run with no diffusion, but especially not recommended with even-order advection

damp_opt=1

- ◆ Upper level diffusive layer
- ◆ Enhanced horizontal and (only for diff_opt=2) vertical diffusion at top
- ◆ Cosine function of height
- ◆ Uses additional parameters
 - zdamp: depth of damping layer
 - dampcoef: nondimensional maximum magnitude of damping
- ◆ Only for idealized cases (for now)

damp_opt=2


- ◆ Upper level relaxation towards 1-d profile
- ◆ Rayleigh (relaxation) layer
- ◆ Cosine function of height
- ◆ Uses additional parameters
 - zdamp: depth of damping layer
 - dampcoef: inverse time scale (s^{-1})
- ◆ Only for idealized cases (for now)

Radiation

Atmospheric temperature
tendency

Surface radiative fluxes

Illustration of Free Atmosphere Radiation Processes



ra_lw_physics=1

RRTM scheme

- ◆ Spectral scheme

- ◆ K-distribution

- ◆ Look-up table fit to accurate calculations

- ◆ Interacts with clouds

- ◆ Ozone/CO₂ from climatology

ra_lw_physics=99

GFDL longwave scheme

- ◆ used in Eta
- ◆ Spectral scheme from global model
- ◆ Also uses tables
- ◆ Interacts with clouds
- ◆ Ozone/CO₂ from climatology

ra_sw_physics=1

MM5 shortwave (Dudhia)

- ◆ Simple downward calculation
- ◆ Clear-sky scattering
- ◆ Water vapor absorption
- ◆ Cloud albedo and absorption

ra_sw_physics=2

Goddard shortwave

- ◆ Spectral method
- ◆ Interacts with clouds
- ◆ Ozone effects

ra_sw_physics=99

GFDL shortwave

- ◆ Used in Eta model
- ◆ Ozone effects
- ◆ Interacts with clouds

radt

Radiation time-step recommendation

- ◆ Radiation is too expensive to call every step
- ◆ Frequency should resolve cloud-cover changes with time
- ◆ $\text{radt} = 1$ minute per km grid size is about right (e.g. $\text{radt} = 10$ for $\text{dx} = 10$ km)

Surface schemes

Surface layer of atmosphere
diagnostics (exchange coeffs)

Soil temperature/moisture/snow,
etc.

sf_sfclay_physics=1

Monin-Obukhov similarity theory

- ◆ Taken from standard relations used in MM5 MRF PBL
- ◆ Provides exchange coefficients to surface (land) scheme
- ◆ Should be used with bl_pbl_physics=1

sf_sfclay_physics=2


Monin-Obukhov similarity theory

- ◆ Modifications due to Janjic

- ◆ Taken from standard relations used in Eta model, including Zilitinkevich thermal roughness length

- ◆ Should be used with bl_pbl_physics=2

Illustration of Surface Processes



sf_surface_physics=1

5-layer thermal diffusion model from MM5

- ◆ Predict ground temp and soil temps
- ◆ Thermal properties depend on land use
- ◆ No effect for water
- ◆ Provides heat and moisture fluxes for PBL

sf_surface_physics=2

Noah Land Surface Model

- ◆ Vegetation effects included
- ◆ Predicts soil temperature and soil moisture in four layers
- ◆ Predicts snow cover and canopy moisture
- ◆ Handles fractional snow cover and frozen soil
- ◆ Diagnoses skin temp and uses emissivity
- ◆ Provides heat and moisture fluxes for PBL

sf_surface_physics=3

RUC Land Surface Model (Smirnova)

- ◆ Vegetation effects included
- ◆ Predicts soil temperature and soil moisture in six layers
- ◆ Multi-layer snow model
- ◆ Provides heat and moisture fluxes for PBL

LANDUSE.TBL

LANDUSE.TBL file (ascii) has land-use properties (vegetation, urban, water, etc.)

- ◆ 24 USGS categories from 30" global dataset
- ◆ Each type is assigned summer/winter value
 - Albedo
 - Emissivity
 - Roughness length
- ◆ Other table properties (thermal inertia, moisture availability, snow albedo effect) are used by 5-layer model
- ◆ Other tables (VEGPARM.TBL, etc.) are used by Noah
- ◆ RUC LSM has internal values

Initializing LSMs

- Noah and RUC LSM require additional fields for initialization
 - Soil temperature
 - Soil moisture
 - Snow liquid equivalent
- Best source is a consistent model-derived dataset
 - Eta/GFS/AGRMET/NNRP for Noah (although some have limited soil levels available)
 - RUC for RUC
- Optimally the resolution, land-use, soil texture, should match the data source model, otherwise there will be a spin-up issue

sst_update=1

Reads lower boundary file periodically to update the sea-surface temperature (otherwise it is fixed with time)


- ◆ For long-period simulations (a week or more)
- ◆ wrflowinp_d01 created by *real*
- ◆ Sea-ice
 - Cannot update sea-ice cover (yet)
 - Treat sea-ice as just cold water (no initial sea ice) if using sst_update
- ◆ Vegetation fraction update can be included in file too

Planetary Boundary Layer

Boundary layer fluxes (heat,
moisture, momentum)

Vertical diffusion

Illustration of PBL Processes



bl_pbl_physics=1

YSU PBL scheme (Hong and Noh)

- ◆ Non-local K mixing in dry convective boundary layer
- ◆ Depth of PBL determined from thermal profile
- ◆ Explicit treatment of entrainment
- ◆ Vertical diffusion depends on Ri in free atmosphere

bl_pbl_physics=2

Mellor-Yamada-Janjic (Eta) PBL

- ◆ 1.5-order, level 2.5, TKE prediction
- ◆ Local K vertical mixing in boundary layer and free atmosphere

bl_pbl_physics=99

MRF PBL scheme (Hong and Pan 1996)

- ◆ Non-local K mixing in dry convective boundary layer
- ◆ Depth of PBL determined from critical Ri number
- ◆ Vertical diffusion depends on Ri in free atmosphere

PBL Scheme Options

PBL schemes can be used for most grid sizes when surface fluxes are present


- ◆ Assumes that PBL eddies are not resolved
- ◆ At grid size $dx \ll 1$ km, this assumption breaks down
- ◆ Can use 3d tke diffusion, but, this is not yet coupled to the actual surface fluxes (future version will have this)
- ◆ Currently 3d tke can only be used with constant specified surface fluxes

Cumulus Parameterization

Atmospheric heat and
moisture/cloud tendencies

Surface rainfall

Illustration of Cumulus Processes



cu_physics=1

New Kain-Fritsch

- ◆ As in MM5 and Eta test version
- ◆ Includes shallow convection
- ◆ CAPE removal time scale
- ◆ Mass flux type with updrafts and downdrafts, entrainment and detrainment
- ◆ Includes cloud detrainment

Cumulus scheme

Recommendations about use

- ◆ For $dx \geq 10$ km: probably need cumulus scheme
- ◆ For $dx \leq 3$ km: probably do not need scheme
 - However, there are cases where the earlier triggering of convection by cumulus schemes help
- ◆ For $dx=3-10$ km, scale separation is a ?
 - No schemes are specifically designed with this range of scales in mind

cu_physics=2

Betts-Miller-Janjic

- ◆ As in Eta model
- ◆ Adjustment type scheme
- ◆ No explicit updraft or downdraft

cu_physics=3

Grell-Devenyi Ensemble

- ◆ Multiple-closure (e.g. CAPE removal, quasi-equilibrium)
- ◆ Multi-parameter (e.g. maximum cap, precipitation efficiency)
- ◆ Explicit updrafts/downdrafts
- ◆ Mean feedback of ensemble is applied
- ◆ Weights can be tuned (spatially, temporally) to optimize scheme (training)

Microphysics


Atmospheric heat and moisture
tendencies

Microphysical rates


Surface rainfall

Illustration of Microphysics Processes


Kessler




WSM3



WSM5



Lin et al./WSM6



mp_physics=1

Kessler scheme

- ◆ Warm rain – no ice
- ◆ Idealized microphysics

mp_physics=2

Purdue Lin et al. scheme

- ◆ 5-class microphysics including graupel
- ◆ Includes ice sedimentation

mp_physics=3

WSM 3-class scheme

- ◆ From Hong, Dudhia and Chen (2004)
- ◆ Replaces NCEP3 scheme
- ◆ 3-class microphysics with ice
- ◆ Ice processes below 0 deg C
- ◆ Ice number is function of ice content
- ◆ Ice sedimentation

mp_physics=4

WSM 5-class scheme

- ◆ Also from Hong, Dudhia and Chen (2004)
- ◆ Replaces NCEP5 scheme
- ◆ 5-class microphysics with ice
- ◆ Supercooled water and snow melt
- ◆ Ice sedimentation

mp_physics=5

Ferrier (current Eta) scheme

- ◆ One prognostic total condensate variable
- ◆ Designed for efficiency
- ◆ Diagnostic ice and water species and liquid fractions

mp_physics=6

WSM 6-class scheme

- ◆ From Hong and Lim (2003 workshop)
- ◆ 6-class microphysics with graupel
- ◆ Ice number concentration as in WSM3 and WSM5
- ◆ Modified accretion

mp_physics=8

Thompson et al. graupel scheme

- ◆ From Thompson et al. (2004, MWR)
- ◆ Newer version of Reisner2 scheme
- ◆ 6-class microphysics with graupel
- ◆ Ice number concentration also predicted (double-moment ice)

mp_physics=98,99

NCEP3,NCEP5

- ◆ Old options from Version 1.3 still available for comparison
- ◆ To be phased out later

mp_zero_out

Microphysics switch (also
mp_zero_out_thresh)

- ◆ 1: all values less than threshold set to zero (except vapor)
- ◆ 2: as 1 but vapor also limited ≥ 0
- ◆ Note: this option will not conserve total water

Microphysics Options


Recommendations about choice

- ◆ Probably not necessary to use a graupel scheme for $dx > 10$ km
 - Updrafts producing graupel not resolved
 - Cheaper scheme may give similar results
- ◆ When resolving individual updrafts, graupel scheme should be used




Physics Interactions

Direct Interactions of Parameterizations






Physics Summary and Plans




Subgrid Turbulence

IN	WORKING ON	PLANNED
<ul style="list-style-type: none">(1) Level 2.5 TKE(2) 3d Smagorinsky(3) Const. coeffs.(4) 2d Smagorinsky (mesoscale)		



Microphysics


IN	WORKING ON	PLANNED
<ul style="list-style-type: none">(1) Kessler(2) Lin et al. [Purdue](3) WSM3(4) WSM5(5) Eta (Ferrier)(6) WSM6(8) Thompson	2-moment schemes (WSM- and Thompson- related)	Goddard



Radiation




	IN	WORKING ON	PLANNED
Long	(1)RRTM (2)Eta (GFDL)	CAM lw	Goddard lw
Short	(1) Dudhia [MM5] (2) Goddard (3) Eta (GFDL)	CAM sw	RRTM sw




Boundary Layer

IN	WORKING ON	PLANNED
(1)YSU (2)M-Y-Janjic (99)MRF	GFS PBL	



Surface

	IN	WORKING ON	PLANNED
Surface layer	(1)MRF Similarity (2)Eta Similarity	GFS surface	
Land surface	(1)5-layer soil temp (2)Noah LSM (3)RUC LSM	CLM	



Cumulus

IN	WORKING ON	PLANNED
<ul style="list-style-type: none">(1) New Kain-Fritsch(2) Betts-Miller-Janjic(3) Grell-Devenyi	<p>Simplified A-S from GFS</p>	



End