

The WRF Preprocessing System: Description of General Functions

Michael Duda



Purpose of this Lecture

In this lecture, our goals are to:


- 1) Understand the purpose of the WPS
 - 2) Learn what each component of the WPS does
-
- The details of *actually running* the WPS are covered in the second WPS lecture
 - *Advanced usage* of the WPS is covered in the third lecture




Purpose of the WPS

The purpose of the WPS is to prepare input to WRF for real-data simulations:

1. Defines simulation domain and ARW nested domains
2. Computes latitude, longitude, map scale factors, Coriolis parameters at every grid point
3. Interpolates time-invariant terrestrial data to simulation gr (e.g., terrain height and soil type)
4. Interpolates time-varying meteorological fields from another model onto simulation domains





WRF Modeling System Flowchart




WPS Program Flowchart

External Data Sources



The *geogrid* program

External Data
Sources




geogrid: think geographical




The *geogrid* program

- For WRF model domains, geogrid defines:
 - Map projection (all domains must use the same)
 - Location of domains
 - Dimensions of domains
- Geogrid provides values for static (time-invariant) fields at each model grid point
 - Compute latitude, longitude, map scale factor, and Coriolis parameters at each grid point
 - Horizontally interpolate static terrestrial data (e.g., topographic height, land use category, soil type, vegetation fraction, monthly surface albedo)







Geogrid: Defining model domains

- First, we choose a map projection to use for the domains; why?
 - The real earth is (roughly) an ellipsoid
 - But WRF computational domains are defined by rectangles in the plane
- NMM uses a rotated latitude-longitude projection
- ARW can use any of the following projections:
 1. Lambert conformal
 2. Mercator
 3. Polar stereographic
 4. Latitude-longitude (for global domain, *must* choose this!)



Supported Projections in ARW



Geogrid: Defining Model Domains

- Define projection of domains using subset of the following parameters
 - **MAP_PROJ**: ‘lambert’, ‘mercator’, ‘polar’, ‘lat-lon’, or ‘rotated_ll’
 - **TRUELAT1**: First true latitude
 - **TRUELAT2**: Second true latitude (*only for Lambert conformal conic projection*)
 - **POLE_LAT**, **POLE_LON**: Location of North Pole in WRF computational grid (*only for ‘lat-lon’*)
 - **STAND_LON**: The meridian parallel to *y*-axis
- All parameters reside in the file *namelist.wps*


*ARW only

See p. 3-9 and 3-34




Geogrid: Defining Model Domains


- Define the area covered (dimensions and location) by coarse domain using the following:
 - **REF_LAT, REF_LON**: The (lat,lon) location of a known location in the domain (*by default, the center point of the domain*)
 - **DX, DY**: Grid distance where map factor = 1
 - For Lambert, Mercator, and polar stereographic: **meters**
 - For (rotated) latitude-longitude: **degrees**
 - **E_WE**: Number of velocity points in west-east direction for ARW; number of mass points in odd rows for NMM
 - **E_SN**: Number of velocity points in south-north direction for ARW; number of rows for NMM




Geogrid: Defining ARW Domains



Geogrid: Defining ARW Domains



Geogrid: Defining NMM Domains




Geogrid: Nesting Basics

- A *nested domain* is a domain that is wholly contained within its *parent domain* and that receives information from its parent, and that may also feed information back to its parent
 - A nested domain has exactly one *parent*
 - A domain may have one or more *children*
- 2-way nests on the same *nesting level* must not overlap in coverage!





Geogrid: Nesting Example

Example configuration – 4 domains




Each domain is assigned a *domain ID #*




Geogrid: Defining Nested ARW Domains

- Define the dimensions and location of nested domains using:
 - **PARENT_ID**: Which domain is the parent?
 - **PARENT_GRID_RATIO**: What is the ratio between grid spacing in parent to grid spacing in this nest?
 - **I_PARENT_START**: *i*-coordinate in parent of this nest's lower-left corner
 - **J_PARENT_START**: *j*-coordinate in parent of this nest's lower-left corner
 - **E_WE**: Number of velocity points in west-east direction
 - **E_SN**: Number of velocity points in south-north direction

See p. 3-15 and 3-33



Geogrid: Defining Nested Domains




The grid spacing of domain 2 is determined by the grid spacing of domain 1 and the `parent_grid_ratio`.

NB: For NMM, the `parent_grid_ratio` is always 3!



Geogrid: Nesting example


Assuming *parent_grid_ratio* = 3



In ARW, nest dimensions must be
($n * \text{parent_grid_ratio} + 1$)
for some integer n


$$112 = 3 * n + 1 \text{ for } n=37$$

$$97 = 3 * n + 1 \text{ for } n=32$$




Geogrid: Interpolating Static Fields


- Given definitions of all computational grids, geogrid interpolates terrestrial, time-invariant fields
 - Topography height
 - Land use categories
 - Soil type (top layer & bottom layer)
 - Annual mean soil temperature
 - Monthly vegetation fraction
 - Monthly surface albedo



Geogrid: Interpolating Static Fields




In general, source data are given on a different projection from the model grid



Geogrid: Interpolation Options

- 4-point bilinear
- 16-point overlapping parabolic
- 4-point average (simple or weighted)
- 16-point average (simple or weighted)
- Grid cell average
- Nearest neighbor
- Breadth-first search

See p. 3-45



Why have so many interpolation options?

- Different interpolators work best for different fields and different relative grid resolutions
 - Some interpolators preserve positive definiteness
 - Some interpolators produce “smoother” fields
 - Some interpolators are best suited for discrete or categorical fields
 - Some are good when going from a fine grid to a coarse g
- Having a choice of how to interpolate fields is good!
 - We'll see in the third WPS lecture how several different options can be used for different regions of the same field



Geogrid: Program Flexibility

- The GEOGRID.TBL file determines
 1. Which fields will be produced by geogrid
 2. What sources of data will be used
 3. How the data will be interpolated/smoothed
 4. Any derived fields (e.g., dominant cat., df/dx)
- Acceptable defaults exist in GEOGRID.TBL, so user will not generally need to edit the file (*but more on this in the third WPS lecture!*)



Geogrid: Program Flexibility

- *geogrid* is flexible enough to ingest and interpolate new static fields
 - handles either continuous or categorical fields
- New data sets must be written to simple binary format
- User needs to add an entry to the file
GEOGRID.TBL



Geogrid: Program Output

- The parameters defining each domain, plus interpolated static fields, are written using the WRF I/O API
 - One file per domain for ARW
 - One file per *nesting level* for NMM
- Filenames: `geo_em.d0n.nc` , or
`geo_nmm.d01.nc`, `geo_nmm_nest.l0k.nc`
(where n is the domain ID # and k is the nest level)
- Example:


| | |
|-----------------------------------|---|
| <code>geo_em.d01.nc</code> | <code>geo_nmm.d01.nc</code> |
| <code>geo_em.d02.nc</code> (nest) | <code>geo_nmm_nest.l01.nc</code> (nest level) |
| <code>geo_em.d03.nc</code> (nest) | <code>geo_nmm_nest.l02.nc</code> (nest level) |




Geogrid: Example Output Fields




LAND-SEA Mask



Topography Height



Top-layer Soil Category




Vegetation Fraction (July)



The *ungrib* program

External Data
Sources




ungrib: think un+grib



The *ungrib* program

- Read GRIB Edition 1 and GRIB Edition 2 files
- Extract meteorological fields
- If necessary, derive required fields from related ones
 - E.g., Compute RH from T, P, and Q
- Write requested fields to an intermediate file format




Ungrib: Vtables

How does ungrib know which fields to extract?


Using Vtables (think: Variable tables)

- Vtables are files that give the GRIB codes for fields to be extracted from GRIB input files
- One Vtable for each source of data
- Vtables are provided for: NAM 104, NAM 212, GFS, AGRMET, and others




Ungrib: Example Vtable

| GRIB1 Param | Level Type | From Level1 | To Level2 | UNGRIB Name | UNGRIB Units | UNGRIB Description |
|----------------|---------------|----------------|--------------|----------------|-----------------|------------------------------------|
| 11 | 100 | * | | T | K | Temperature |
| 33 | 100 | * | | U | m s-1 | U |
| 34 | 100 | * | | V | m s-1 | V |
| 52 | 100 | * | | RH | % | Relative Humidity |
| 7 | 100 | * | | HGT | m | Height |
| 11 | 105 | 2 | | T | K | Temperature at 2 m |
| 52 | 105 | 2 | | RH | % | Relative Humidity at 2 m |
| 33 | 105 | 10 | | U | m s-1 | U at 10 m |
| 34 | 105 | 10 | | V | m s-1 | V at 10 m |
| 1 | 1 | 0 | | PSFC | Pa | Surface Pressure |
| 130 | 102 | 0 | | PMSL | Pa | Sea-level Pressure |
| 144 | 112 | 0 | 10 | SM000010 | kg m-3 | Soil Moist 0-10 cm below grn layer |
| 144 | 112 | 10 | 40 | SM010040 | kg m-3 | Soil Moist 10-40 cm below grn laye |
| 144 | 112 | 40 | 100 | SM040100 | kg m-3 | Soil Moist 40-100 cm below grn lay |
| 144 | 112 | 100 | 200 | SM100200 | kg m-3 | Soil Moist 100-200 cm below gr lay |
| 85 | 112 | 0 | 10 | ST000010 | K | T 0-10 cm below ground layer (Uppe |
| 85 | 112 | 10 | 40 | ST010040 | K | T 10-40 cm below ground layer (Upp |
| 85 | 112 | 40 | 100 | ST040100 | K | T 40-100 cm below ground layer (Up |
| 85 | 112 | 100 | 200 | ST100200 | K | T 100-200 cm below ground layer (E |
| 91 | 1 | 0 | | SEAICE | propnrt | Ice flag |
| 81 | 1 | 0 | | LANDSEA | propnrt | Land/Sea flag (1=land,2=sea in GRI |
| 7 | 1 | 0 | | HGT | m | Terrain field of source analysis |
| 11 | 1 | 0 | | SKINTEMP | K | Skin temperature (can use for SST |
| 65 | 1 | 0 | | SNOW | kg m-2 | Water equivalent snow depth |
| 223 | 1 | 0 | | CANWAT | kg m-2 | Plant Canopy Surface Water |
| 224 | 1 | 0 | | SOILCAT | Tab4.213 | Dominant soil type category |
| 225 | 1 | 0 | | VEGCAT | Tab4.212 | Dominant land use category |



Ungrib: GRIB2 Vtable Entries

| | GRIB2 | GRIB2 | GRIB2 | GRIB2 |
|---|-------|-------|-------|-------|
| | Discp | Catgy | Param | Level |
| metgrid | | | | |
| Description | | | | |
| Temperature | 0 | 0 | 0 | 100 |
| U | 0 | 2 | 2 | 100 |
| V | 0 | 2 | 3 | 100 |
| Relative Humidity | 0 | 1 | 1 | 100 |
| Height | 0 | 3 | 5 | 100 |
| Temperature at 2 m | 0 | 0 | 0 | 103 |
| Relative Humidity at 2 m | 0 | 1 | 1 | 103 |
| U at 10 m | 0 | 2 | 2 | 103 |
| V at 10 m | 0 | 2 | 3 | 103 |
| Surface Pressure | 0 | 3 | 0 | 1 |
| Sea-level Pressure | 0 | 3 | 1 | 101 |
| Soil Moist 0-10 cm below grn layer (Up) | 2 | 0 | 192 | 106 |
| Soil Moist 10-40 cm below grn layer | 2 | 0 | 192 | 106 |
| Soil Moist 40-100 cm below grn layer | 2 | 0 | 192 | 106 |
| Soil Moist 100-200 cm below gr layer | 2 | 0 | 192 | 106 |
| Soil Moist 10-200 cm below gr layer | 2 | 0 | 192 | 106 |
| T 0-10 cm below ground layer (Upper) | 0 | 0 | 0 | 106 |
| T 10-40 cm below ground layer (Upper) | 0 | 0 | 0 | 106 |
| T 40-100 cm below ground layer (Upper) | 0 | 0 | 0 | 106 |
| T 100-200 cm below ground layer (Bottom) | 0 | 0 | 0 | 106 |
| T 10-200 cm below ground layer (Bottom) | 0 | 0 | 0 | 106 |
| Ice flag | 0 | 2 | 0 | 1 |
| Land/Sea flag (1=land, 0 or 2=sea) | 2 | 0 | 0 | 1 |
| Terrain field of source analysis | 2 | 0 | 7 | 1 |
| Skin temperature (can use for SST also) | 0 | 0 | 0 | 1 |
| Water equivalent snow depth | 0 | 1 | 13 | 1 |
| Dominant soil type cat. (not in GFS file) | 2 | 3 | 0 | 1 |
| Dominant land use cat. (not in GFS file) | 2 | 0 | 198 | 1 |



Ungrib: Vtables

What if a data source has no existing Vtable?

Create a Vtable


- Get a listing of GRIB codes for fields in the source
 - Check documentation from originating center or use utility such as *wgrib*, *g1print*, *g2print*
- Use existing Vtable as a template
- Check documentation in Chapter 3 of the Users' Guide for more information about Vtables

See p. 3-27



Ungrib: Intermediate File Format

- After extracting fields listed in Vtable, ungrib writes those fields to intermediate format
- For meteorological data sets not in GRIB format, the user may write to intermediate format directly
 - Allows WPS to ingest new data sources; basic programming required of user See p. 3-25
 - Simple intermediate file format is easily read/written using routines from WPS ([read_met_module.F](#) and [write_met_module.F](#))



Ungrib: Program Output

- Output files named *FILE:YYYY-MM-DD_HH*
 - *YYYY* is year of data in the file; *MM* is month; *DD* day; *HH* is hour
 - All times are UTC
- Example:
 - FILE:2007-07-24_00*
 - FILE:2007-07-24_06*
 - FILE:2007-07-24_12*

ungrib can also write intermediate files in MM5 or WRF SI format! (*To allow for use of GRIB2 data in MM5, for example*)




Ungrib: Obtaining GRIB Data

- Where does one get GRIB data?
 - User's responsibility
 - Some free data are available from NCAR and NCEP. See
 - <http://www.mmm.ucar.edu/wrf/users/>
 - > under the “Downloads” tab:
 - Some NCEP data in the past year
 - NCEP operational data available daily



The *metgrid* program

External Data
Sources



metgrid: think meteorological




The *metgrid* program

- Horizontally interpolate meteorological data (*extracted by ungrib*) to simulation domains (*defined by geogrid*)
 - Masked interpolation for masked fields
- Rotate winds to WRF grid
 - i.e., rotate so that U-component is parallel to x-axis
V-component is parallel to y-axis



Metgrid: ARW Grid Staggering

- For ARW, wind U-component interpolated to “u” staggering
- Wind V-component interpolated to “v” staggering
- Other meteorological fields interpolated to “ θ ” staggering by default (*can change this!*)




A single ARW cell, with “u”, “ θ ” and “v” points labeled.



Metgrid: NMM Grid Staggering

- For NMM, wind U- and V-components interpolated to “V” staggering
- Other meteorological fields interpolated to “H” staggering by default (*can change this!*)



An NMM grid showing “V”, and “H” points.



Metgrid: Interpolation Options*

- 4-point bilinear
- 16-point overlapping parabolic
- 4-point average (simple or weighted)
- 16-point average (simple or weighted)
- Grid cell average
- Nearest neighbor
- Breadth-first search

* These are the same options available for geogrid!




Metgrid: Masked Interpolation

- *Masked fields* may only have valid data at a subset of grid points
 - E.g.: SST field only valid on water points
- When metgrid interpolates masked fields, it must know which points are invalid (masked)
 - Can use separate mask field (e.g., LANDSEA)
 - Can rely on special values (e.g., 1×10^{30}) in field itself to identify masked grid points



Metgrid: Masked Interpolation




Suppose we need to interpolate to point X

- Using **red** points as valid data can give a bad interpolated value!
- Masked interpolation only uses valid **blue** points to interpolate to X


*Not every interpolation option can handle masked points;
we'll address this issue in the third lecture*




Example: Masked Interpolation



Skin temperature field interpolated from GFS 0.5-deg field with no mask using a sixteen-point interpolator.



Skin temperature field interpolated using masks: GFS water points interpolated to model water points; GFS land points interpolated to model land points.




Metgrid: Wind Rotation


- Input wind fields (U-component + V-component) are either:
 - **Earth-relative**: U-component = westerly component; V-component = southerly component
 - **Relative to source grid**: U-component (V-component) parallel to source model x-axis (y-axis)
- WRF expects wind components to be relative to the simulation grid



Metgrid: Wind Rotation Example



A wind vector, shown in terms of its U and V components with respect to the source grid.




The same vector, in terms of its U and V components with respect to the WRF simulation grid.

This process may require *two* rotations: one from source grid to earth grid and a second from earth grid to WRF grid




Metgrid: Constant Fields

- For short simulations, some fields may be constant
 - E.g., SST or sea-ice fraction
- Use namelist option `CONSTANTS_NAME` option to specify such fields:
 - `CONSTANTS_NAME = 'SST_FILE:2007-07-24_00'`



Metgrid: Program Flexibility

- *metgrid* is capable of interpolating both isobar and native vertical coordinate data sets
- User may specify interpolation methods and related options in the **METGRID.TBL** file
 - **METGRID.TBL** file similar in format to the file **GEOGRID.TBL**




Metgrid: Program Output




- For coarse domain, one file per time period
 - In ARW, we also get the first time period for all nested grids
- Files contain static fields from geogrid plus interpolated meteorological fields
- Filenames:

ARW: `met_em.d0n.YYYY-MM-DD_HH:mm:ss.nc`
(where n is the domain ID #)


NMM: `met_nmm.d01.YYYY-MM-DD_HH:mm:ss.nc`




Metgrid: Example Output



Mean Sea-level Pressure



WPS Summary



And finally...

Vertical interpolation to WRF eta levels is performed in the *real* or *real_nmm* program

