



# WRF Physics Options

Jimmy Dudhia

# WRF Physics

- ◆ Turbulence/Diffusion (diff\_opt, km\_opt)
- ◆ Radiation
  - Longwave (ra\_lw\_physics)
  - Shortwave (ra\_sw\_physics)
- ◆ Surface
  - Surface layer (sf\_sfclay\_physics)
  - Land/water surface (sf\_surface\_physics)
- ◆ PBL (bl\_physics)
- ◆ Cumulus parameterization (cu\_physics)
- ◆ Microphysics (mp\_physics)

# Turbulence/Diffusion


Sub-grid eddy mixing effects on

all fields, e.g.  $\frac{\partial}{\partial x} K_h \frac{\partial}{\partial x} \theta + \frac{\partial}{\partial y} K_h \frac{\partial}{\partial y} \theta + \frac{\partial}{\partial z} K_v \frac{\partial}{\partial z} \theta$

# diff\_opt=1

- ◆ 2<sup>nd</sup> order diffusion on model levels
  - Constant vertical coefficient (kvdif) or use with PBL
  - For theta, only perturbation from base state is diffused
- ◆ km\_opt selects method to compute K
  - 1: constant (khdif and kvdif used)
  - 4: 2D Smagorinsky (deformation based on horizontal wind for horizontal diffusion only)

# Difference between diff\_opt 1 and 2




diff\_opt=1

Horizontal diffusion acts along model levels

Simpler numerical method with only neighboring points on the same model level

# Difference between diff\_opt 1 and 2



diff\_opt=2

Horizontal diffusion acts along model levels

Numerical method includes vertical correction term  
using more grid points

# diff\_opt=2

- ◆ 2<sup>nd</sup> order horizontal diffusion
- ◆ Allows for terrain-following coordinate
- ◆ km\_opt selects method to compute K
  - 1: constant (khdif and kvdif used)
  - 2: 1.5-order TKE prediction
  - 3: Smagorinsky (deformation/stability based K)
  - 4: 2D Smagorinsky (deformation based on horizontal wind for horizontal diffusion only)

# diff\_opt=2 (continued)

- ◆ mix\_full\_fields=.true.: vertical diffusion acts on full (not perturbation) fields (recommended, but default = .false.)
- ◆ mix\_isotropic=1: same length scale used for horizontal and vertical diffusion (for  $dx \approx dz$ )
- ◆ Idealized constant surface fluxes can be added in diff\_opt=2 using namelist (dynamics section). Not available for diff\_opt=1.
  - tke\_drag\_coefficient ( $C_D$ )
  - tke\_heat\_flux ( $=H/\rho c_p$ )
  - Must use isfflx=0 to use these switches



# diff\_opt=2 (continued)

- ◆ Explicit large-eddy simulation (LES) PBL in real-data cases (V3) or idealized cases
  - bl\_pbl\_physics = 0
    - ◆ isfflx = 0 (idealized drag and heat flux from namelist)
    - ◆ isfflx = 1 (drag and heat flux from physics)
      - sf\_sfclay\_physics=1
      - sf\_surface\_physics (choose non-zero option)
    - ◆ isfflx = 2 (drag from physics, heat flux from tke\_heat\_flux)
      - sf\_sfclay\_physics=1
  - km\_opt = 2 or 3
  - mix\_isotropic=1 (if dx and dz are of same order)
- ◆ Not available for diff\_opt=1.

# Diffusion Option Choice

- ◆ Real-data case with PBL physics on
  - Best is diff\_opt=1, km\_opt=4
  - This complements vertical diffusion done by PBL scheme
- ◆ High-resolution real-data cases (~100 m grid)
  - No PBL
  - diff\_opt=2; km\_opt=2,3 (tke or Smagorinsky scheme)
- ◆ idealized cloud-resolving modeling (smooth or no topography)
  - diff\_opt=2; km\_opt=2,3
- ◆ Complex topography with no PBL scheme
  - diff\_opt=2 is more accurate for sloped coordinate surfaces, and prevents diffusion up/down valley sides
- ◆ Note: WRF can run with no diffusion (diff\_opt=0)

# diff\_6th\_opt

- ◆ 6<sup>th</sup> order optional added horizontal diffusion on model levels
  - Used as a numerical filter for  $2 \times dx$  noise
  - Suitable for idealized and real-data cases
  - Affects all advected variables including scalars
- ◆ diff\_6th\_opt
  - 0: none (default)
  - 1: on (can produce negative water)
  - 2: on and prohibit up-gradient diffusion (better for water conservation)
- ◆ diff\_6th\_factor
  - Non-dimensional strength (typical value 0.12, 1.0 corresponds to complete removal of  $2 \times dx$  wave in a time-step)

# damp\_opt=1

- ◆ Upper level diffusive layer
- ◆ Enhanced horizontal diffusion at top
- ◆ Also enhanced vertical diffusion at top for diff\_opt=2
- ◆ Cosine function of height
- ◆ Uses additional parameters
  - zdamp: depth of damping layer
  - dampcoef: nondimensional maximum magnitude of damping
- ◆ Works for idealized cases and real-data since 2.2 release

# damp\_opt=2

- ◆ Upper level relaxation towards 1-d profile
- ◆ Rayleigh (relaxation) layer
- ◆ Cosine function of height
- ◆ Uses additional parameters
  - zdamp: depth of damping layer
  - dampcoef: inverse time scale ( $s^{-1}$ )
- ◆ Works for idealized cases only

# damp\_opt=3

- ◆ “W-Rayleigh” (relaxation) layer
- ◆ Upper level relaxation towards zero vertical motion
- ◆ Cosine function of height
- ◆ Uses additional parameters
  - zdamp: depth of damping layer
  - dampcoef: inverse time scale ( $s^{-1}$ )
- ◆ Works for idealized and real-data cases
- ◆ Applied in small time-steps (dampcoef=0.2 is stable)




# Radiation

Atmospheric temperature  
tendency

Surface radiative fluxes

## Illustration of Free Atmosphere Radiation Processes





# ra\_lw\_physics=1

RRTM scheme

- ◆ Spectral scheme
- ◆ K-distribution
- ◆ Look-up table fit to accurate calculations
- ◆ Interacts with clouds (1/0 fraction)
- ◆ Ozone profile specified
- ◆ CO2 constant (well-mixed)

# ra\_lw\_physics=3

CAM3 scheme

- ◆ Spectral scheme
- ◆ 8 longwave bands
- ◆ Look-up table fit to accurate calculations
- ◆ Interacts with clouds
- ◆ Can interact with trace gases and aerosols
- ◆ Ozone profile function of month, latitude
- ◆ CO2 changes based on year (new in V3.1)

# ra\_lw\_physics=4

RRTMG longwave scheme (New in V3.1)

- ◆ Spectral scheme
- ◆ 16 longwave bands (K-distribution)
- ◆ Look-up table fit to accurate calculations
- ◆ Interacts with clouds (MCICA method)
- ◆ Can interact with trace gases and aerosols
- ◆ Ozone profile specified
- ◆ CO2 and trace gases specified

# ra\_lw\_physics=99

GFDL longwave scheme

- ◆ used in Eta/NMM
- ◆ Default code is used with Ferrier microphysics
  - Remove #define to compile for use without Ferrier
- ◆ Spectral scheme from global model
- ◆ Also uses tables
- ◆ Interacts with clouds (cloud fraction)
- ◆ Ozone profile based on season, latitude
- ◆ CO2 fixed

# ra\_sw\_physics=1

MM5 shortwave (Dudhia)

- ◆ Simple downward calculation
- ◆ Clear-sky scattering
  - swrad\_scattuning parameter
    - ◆ 1.0 = 10% scattered, 0.5=5%, etc.
  - WRF-Chem aerosol effect (PM2.5)
- ◆ Water vapor absorption
- ◆ Cloud albedo and absorption
- ◆ No ozone effect (model top below 50 hPa OK)
- ◆ Surface slope and shading switches (slope\_rad, topo\_shading)

# ra\_sw\_physics=2

Goddard shortwave

- ◆ Spectral method
- ◆ Interacts with clouds
- ◆ Ozone profile (tropical, summer/winter, mid-lat, polar)
- ◆ CO2 fixed
- ◆ WRF-Chem optical thicknesses

# ra\_sw\_physics=3

## CAM3 shortwave

- ◆ Spectral method (19 bands)
- ◆ Interacts with clouds
- ◆ Ozone/CO2 profile as in CAM longwave
- ◆ Can interact with aerosols and trace gases
- ◆ Note: CAM schemes need some extra namelist items (see README.namelist)

# ra\_sw\_physics=4

RRTMG shortwave (New in V3.1)

- ◆ Spectral method (14 bands)
- ◆ Interacts with clouds (MCICA method)
- ◆ Ozone/CO2 profile as in RRTMG longwave
- ◆ Can interact with aerosols
- ◆ Trace gases specified



# ra\_sw\_physics=99

GFDL shortwave

- ◆ Used in Eta/NMM model
- ◆ Default code is used with Ferrier microphysics (see GFDL longwave)
- ◆ Ozone/CO2 profile as in GFDL longwave
- ◆ Interacts with clouds (and cloud fraction)

# radt

## Radiation time-step recommendation

- ◆ Radiation is too expensive to call every step
- ◆ Frequency should resolve cloud-cover changes with time
- ◆  $\text{radt} = 1$  minute per km grid size is about right (e.g.  $\text{radt} = 10$  for  $\text{dx} = 10$  km)
- ◆ Each domain can have its own value but recommend using same value on all 2-way nests

# nrads/nradl

## Radiation time-step recommendation


- Number of fundamental steps per radiation call
- Operational setting should be 3600/dt
- Higher resolution could be used, e.g. 1800/dt
- Recommend same value for all nested domains




# Surface schemes

Surface layer of atmosphere  
diagnostics (exchange/transfer  
coeffs)

Land Surface: Soil temperature  
/moisture /snow prediction /sea-  
ice temperature



# Surface Physics Components



# Surface Fluxes

◆ Heat, moisture and momentum

$$H = \rho c_p u_* \theta_* \quad E = \rho u_* q_* \quad \tau = \rho u_* u_*$$

$$u_* = \frac{kV_r}{\ln(z_r / z_0) - \psi_m} \quad \theta_* = \frac{k\Delta\theta}{\ln(z_r / z_{0h}) - \psi_h} \quad q_* = \frac{k\Delta q}{\ln(z_r / z_{0q}) - \psi_h}$$

Subscript  $r$  is reference level (lowest model level, or 2 m or 10 m)  
 $z_0$  are the roughness lengths

# Roughness Lengths

- ◆ Roughness lengths are a measure of the “initial” length scale of surface eddies, and generally differ for velocity and scalars
- ◆ Roughness length depends on land-use type
- ◆ Some schemes use smaller roughness length for heat than for momentum
- ◆ For water points roughness length is a function of surface wind speed

# Exchange Coefficient

◆  $C_{hs}$  is the exchange coefficient for heat, defined such that

$$H = \rho c_p C_{hs} \Delta \theta$$

It is related to the roughness length and  $u^*$  by

$$C_{hs} = \frac{ku_*}{\ln\left(\frac{z}{z_0}\right) - \psi_h}$$



# sf\_sfclay\_physics=1

Monin-Obukhov similarity theory

- ◆ Taken from standard relations used in MM5 MRF PBL
- ◆ Provides exchange coefficients to surface (land) scheme
- ◆ Should be used with bl\_pbl\_physics=1 or 99

# sf\_sfclay\_physics=2

Monin-Obukhov similarity theory

- ◆ Modifications due to Janjic
- ◆ Taken from standard relations used in NMM model, including Zilitinkevich thermal roughness length
- ◆ Should be used with bl\_pbl\_physics=2

NMM only

# sf\_sfclay\_physics=3

GFS Monin-Obukhov similarity theory

- ◆ For use with NMM-LSM
- ◆ Should be used with bl\_pbl\_physics=3

# sf\_sfclay\_physics=4

QNSE Monin-Obukhov similarity theory  
(New in V3.1)

- ◆ For use with QNSE-PBL
- ◆ Should be used with bl\_pbl\_physics=4
- ◆ Very similar to MYJ SFC
- ◆ New stability functions

# sf\_sfclay\_physics=5

MYNN Monin-Obukhov similarity theory  
(New in V3.1)


- ◆ For use with MYNN-PBL
- ◆ Should be used with bl\_pbl\_physics=5

# sf\_sfclay\_physics=7

Pleim-Xiu surface layer (EPA)

- ◆ For use with PX LSM and ACM PBL
  - Should be used with sf\_surface\_physics=7 and bl\_pbl\_physics=7
- ◆ New in Version 3

## Illustration of Surface Processes



# sf\_surface\_physics=1

5-layer thermal diffusion model from MM5

- ◆ Predict ground temp and soil temps
- ◆ Thermal properties depend on land use
- ◆ No effect for water (Version 3 has ocean mixed-layer model for hurricane applications)
- ◆ No soil moisture or snow-cover prediction
- ◆ Moisture availability based on land-use only
- ◆ Provides heat and moisture fluxes for PBL
- ◆ May be available for NMM in Version 3



# sf\_surface\_physics=2

Noah Land Surface Model (Unified ARW/NMM version in Version 3)

- ◆ Vegetation effects included
- ◆ Predicts soil temperature and soil moisture in four layers and diagnoses skin temperature
- ◆ Predicts snow cover and canopy moisture
- ◆ Handles fractional snow cover and frozen soil
- ◆ New time-varying snow albedo (in V3.1)
- ◆ Provides heat and moisture fluxes for PBL
- ◆ Noah has 2 Urban Canopy Model options (sf\_urban\_physics, ARW only)

# sf\_urban\_physics=1

Urban Canopy Model (UCM, Kusaka et al.)

- ◆ Sub-grid wall, roof, and road effects on radiation and fluxes
- ◆ Anthropogenic heat source can be specified
- ◆ Can use low, medium and high density urban categories

# sf\_urban\_physics=2

Building Environment Parameterization (BEP, Martilli et al.)

- ◆ Sub-grid wall, roof, and road effects on radiation and fluxes
- ◆ Can be used with MYJ PBL or BouLac PBL to represent buildings higher than lowest model levels (Multi-layer urban model)
- ◆ Needs additional sub-grid building fractional area information

# sf\_surface\_physics=3

## RUC Land Surface Model (Smirnova)

- ◆ Vegetation effects included
- ◆ Predicts soil temperature and soil moisture in six layers
- ◆ Multi-layer snow model
- ◆ Provides heat and moisture fluxes for PBL

# sf\_surface\_physics=7

## Pleim-Xiu Land Surface Model (EPA)

- ◆ New in Version 3
- ◆ Vegetation effects included
- ◆ Predicts soil temperature and soil moisture in two layers
- ◆ Simple snow-cover model
- ◆ Provides heat and moisture fluxes for PBL

# LANDUSE.TBL

Text (ASCII) file that has land-use properties for slab model (vegetation, urban, water, etc.)

- ◆ From Version 3.1 Noah LSM does not use this table
- ◆ 24 USGS categories or 20 MODIS categories (new) from 30" global dataset
- ◆ Each type is assigned summer/winter value
  - Albedo
  - Emissivity
  - Roughness length
- ◆ Other table properties (thermal inertia, moisture availability, snow albedo effect) are used by 5-layer model
- ◆ Also note
  - Other tables (VEGPARM.TBL, etc.) are used by Noah
  - RUC LSM uses same table files after Version 3

# Initializing LSMs

- Noah and RUC LSM require additional fields for initialization
  - Soil temperature
  - Soil moisture
  - Snow liquid equivalent
- These are in the Grib files, but are not from observations
- They come from “offline” models driven by observations (rainfall, radiation, surface temperature, humidity wind)

# Initializing LSMs

- There are consistent model-derived datasets for Noah and RUC LSMs
  - Eta/GFS/AGRMET/NNRP for Noah (although some have limited soil levels available)
  - RUC for RUC
- But, resolution of mesoscale land-use means there will be inconsistency in elevation, soil type and vegetation
- This leads to spin-up as adjustments occur in soil temperature and moisture
- This spin-up can only be avoided by running offline model on the same grid (e.g. HRLDAS for Noah)
- Cycling land state between forecasts also helps, but may propagate errors (e.g in rainfall effect on soil moisture)



# sst\_update=1

Reads lower boundary file periodically to update the sea-surface temperature (otherwise it is fixed with time)

- ◆ For long-period simulations (a week or more)
- ◆ wrflowinp\_d0n created by *real*
- ◆ Sea-ice can be updated since Version 3.0
- ◆ Vegetation fraction update is included
  - Allows seasonal change in albedo, emissivity, roughness length in Noah LSM
- ◆ usemonalb=.true. to use monthly albedo input

# Regional Climate Options

- ◆ New in V3.1
- ◆ `tmn_update=1` - updates deep-soil temperature for multi-year future-climate runs
- ◆ `sst_skin=1` - adds diurnal cycle to sea-surface temperature
- ◆ `bucket_mm` and `bucket_J` - a more accurate way to accumulate water and energy for long-run budgets
- ◆ No-leap-year compilation option for CCSM-driven runs

# Hurricane Options

- ◆ Ocean Mixed Layer Model (omlcall=1)
  - Use with sf\_surface\_physics=1
  - 1-d slab ocean mixed layer (specified initial depth)
  - Includes wind-driven ocean mixing for SST cooling feedback
- ◆ Alternative surface-layer option for high-wind ocean surface (isftcflx=1)
  - Use with sf\_sfclay\_physics=1
  - Modifies Charnock relation to give less surface friction at high winds (lower Cd)
  - Modifies surface enthalpy (Ck, heat/moisture) formulation

# Fractional Sea Ice

- ◆ fractional\_seaice=1 - with input sea-ice fraction data can partition land/water fluxes within a grid box
- ◆ New in Version 3.1


# Planetary Boundary Layer

Boundary layer fluxes (heat,  
moisture, momentum)

Vertical diffusion

$$\frac{\partial}{\partial z} K_v \frac{\partial}{\partial z} \theta$$

## Illustration of PBL Processes



# bl\_pbl\_physics=1

YSU PBL scheme (Hong, Noh and Dudhia 2006)

- ◆ Parabolic K profile mixing in dry convective boundary layer
- ◆ Troen-Mahrt countergradient flux (non-local)

$$\frac{\partial}{\partial z} \left( K_v \frac{\partial}{\partial z} \theta + \Gamma \right)$$

- ◆ Depth of PBL determined from thermal profile
- ◆ Explicit treatment of entrainment
- ◆ Vertical diffusion depends on Ri in free atmosphere
- ◆ New stable surface BL mixing using bulk Ri

# bl\_pbl\_physics=2

Mellor-Yamada-Janjic (Eta/NMM) PBL

- ◆ 1.5-order, level 2.5, TKE prediction
- ◆ Local TKE-based vertical mixing in boundary layer and free atmosphere
- ◆ TKE\_MYJ is advected by NMM, not by ARW (yet)



# bl\_pbl\_physics=4

QNSE (Quasi-Normal Scale Elimination)  
PBL from Galperin and Sukoriansky

- ◆ 1.5-order, level 2.5, TKE prediction
- ◆ Local TKE-based vertical mixing in boundary layer and free atmosphere
- ◆ New theory for stably stratified case
- ◆ New in V3.1

# bl\_pbl\_physics=5 and 6

MYNN (Nakanishi and Niino) PBL

- ◆ (5) 1.5-order, level 2.5, TKE prediction, OR
- ◆ (6) 2nd-order, level 3, TKE,  $\theta'^2$ ,  $q'^2$  and  $\theta'q'$  prediction
- ◆ Local TKE-based vertical mixing in boundary layer and free atmosphere
- ◆ New in V3.1

# bl\_pbl\_physics=3

## GFS PBL

- ◆ 1st order Troen-Mahrt
- ◆ Closely related to MRF PBL
- ◆ Non-local-K vertical mixing in boundary layer and free atmosphere

# bl\_pbl\_physics=7

Asymmetrical Convective Model, Version 2  
(ACM2) PBL (Pleim and Chang)

- ◆ Blackadar-type thermal mixing upwards from surface layer
- ◆ Local mixing downwards
- ◆ PBL height from critical bulk Richardson number

# bl\_pbl\_physics=8

BouLac PBL (Bougeault and Lacarrère)

- ◆ TKE prediction scheme
- ◆ Designed to work with multi-layer urban model (BEP)
- ◆ New in V3.1

# bl\_pbl\_physics=99

MRF PBL scheme (Hong and Pan 1996)

- ◆ Non-local-K mixing in dry convective boundary layer
- ◆ Depth of PBL determined from critical Ri number
- ◆ Vertical diffusion depends on Ri in free atmosphere

# bldt

- ◆ Minutes between boundary layer/LSM calls
- ◆ Typical value is 0 (every step)

# nphs

- ◆ Time steps between PBL/turbulence/LSM calls
- ◆ Typical value is chosen to give a frequency of 1-3 minutes, i.e.  $60/dt$  to  $180/dt$
- ◆ Also used for microphysics



# PBL Scheme Options

PBL schemes can be used for most grid sizes when surface fluxes are present

- ◆ With YSU, ACM2, GFS and MRF PBL schemes, lowest full level should be .99 or .995 (not too close to 1)
- ◆ TKE schemes can use thinner surface layers
- ◆ Assumes that PBL eddies are not resolved
- ◆ At grid size  $dx \ll 1$  km, this assumption breaks down
  - Can use 3d diffusion instead of a PBL scheme in Version 3 (coupled to surface physics)
  - Works best when  $dx$  and  $dz$  are comparable

# diff\_opt=2 (repeated)

- ◆ Explicit large-eddy simulation (LES) PBL in real-data cases (V3) or idealized cases
  - bl\_pbl\_physics = 0
    - ◆ isfflx = 0 (idealized drag and heat flux from namelist)
    - ◆ isfflx = 1 (drag and heat flux from physics)
      - sf\_sfclay\_physics=1
      - sf\_surface\_physics (choose non-zero option)
    - ◆ isfflx = 2 (drag from physics, heat flux from tke\_heat\_flux)
      - sf\_sfclay\_physics=1
  - km\_opt = 2 or 3
  - mix\_isotropic=1 (if dx and dz are of same order)
- ◆ Not available for diff\_opt=1.

# Gravity Wave Drag

(gwd\_opt=1 for ARW, 2 for NMM)


- ◆ ARW scheme from Hong et al. New in V3.1
- ◆ Accounts for orographic gravity wave effect on momentum profile
- ◆ Extra sub-grid orographic information comes from geogrid
- ◆ Probably needed only if all below apply
  - $dx > 10$  km
  - Simulations longer than 5 days
  - Domains including mountains



# Cumulus Parameterization

Atmospheric heat and  
moisture/cloud tendencies  
Surface rainfall

## Illustration of Cumulus Processes



# cu\_physics=1

## New Kain-Fritsch

- ◆ As in MM5 and Eta/NMM test version
- ◆ Includes shallow convection (no downdrafts)
- ◆ Low-level vertical motion in trigger function
- ◆ CAPE removal time scale closure
- ◆ Mass flux type with updrafts and downdrafts, entrainment and detrainment
- ◆ Includes cloud, rain, ice, snow detrainment
- ◆ Clouds persist over convective time scale (recalculated every convective step in NMM)
- ◆ Old KF is option 99

# cu\_physics=2

## Betts-Miller-Janjic

- ◆ As in NMM model (Janjic 1994)
- ◆ Adjustment type scheme
- ◆ Deep and shallow profiles
- ◆ BM saturated profile modified by cloud efficiency, so post-convective profile can be unsaturated in BMJ
- ◆ No explicit updraft or downdraft
- ◆ No cloud detrainment
- ◆ Scheme changed significantly since V2.1

# cu\_physics=3

## Grell-Devenyi Ensemble

- ◆ Multiple-closure (CAPE removal, quasi-equilibrium, moisture convergence, cloud-base ascent) - 16 mass flux closures
- ◆ Multi-parameter (maximum cap, precipitation efficiency) - e.g. 3 cap strengths, 3 efficiencies
- ◆ Explicit updrafts/downdrafts
- ◆ Includes cloud and ice detrainment
- ◆ Mean feedback of ensemble is applied
- ◆ Weights can be tuned (spatially, temporally) to optimize scheme (training)



# cu\_physics=4

Simplified Arakawa-Schubert (SAS) GFS scheme

- ◆ Quasi-equilibrium scheme
- ◆ Related to Grell scheme in MM5
- ◆ Includes cloud and ice detrainment
- ◆ Downdrafts and single, simple cloud

# cu\_physics=5

## Grell-3d

- ◆ As GD, but slightly different ensemble
- ◆ Includes cloud and ice detrainment
- ◆ Subsidence is spread to neighboring columns
  - This makes it more suitable for  $< 10$  km grid size than other options
  - `cugd_avgdx=1` (default), 3(spread subsidence)
- ◆ Mean feedback of ensemble is applied
- ◆ Weights can be tuned (spatially, temporally) to optimize scheme (training)

# cutd

- ◆ Time steps between cumulus scheme calls
- ◆ Typical value is 5 minutes

## ncnvc

- ◆ Time steps between cumulus parameterization calls
- ◆ Typically 10 - same as NPHS

# Cumulus scheme

## Recommendations about use

- ◆ For  $dx \geq 10$  km: probably need cumulus scheme
- ◆ For  $dx \leq 3$  km: probably do not need scheme
  - However, there are cases where the earlier triggering of convection by cumulus schemes help
- ◆ For  $dx=3-10$  km, scale separation is a question
  - No schemes are specifically designed with this range of scales in mind
- ◆ Issues with 2-way nesting when physics differs across nest boundaries (seen in precip field on parent domain)
  - best to use same physics in both domains or 1-way nesting



# Microphysics


Atmospheric heat and moisture tendencies

Microphysical rates


Surface rainfall

# Illustration of Microphysics Processes


Kessler




WSM3




Ferrier



WSM5



Lin et al./WSM6



# mp\_physics=1

Kessler scheme

- ◆ Warm rain – no ice
- ◆ Idealized microphysics
- ◆ Time-split rainfall



# mp\_physics=2

Purdue Lin et al. scheme

- ◆ 5-class microphysics including graupel
- ◆ Includes ice sedimentation and time-split fall terms

# mp\_physics=3

WSM 3-class scheme

- ◆ From Hong, Dudhia and Chen (2004)
- ◆ Replaces NCEP3 scheme
- ◆ 3-class microphysics with ice
- ◆ Ice processes below 0 deg C
- ◆ Ice number is function of ice content
- ◆ Ice sedimentation and time-split fall terms

# mp\_physics=4

## WSM 5-class scheme

- ◆ Also from Hong, Dudhia and Chen (2004)
- ◆ Replaces NCEP5 scheme
- ◆ 5-class microphysics with ice
- ◆ Supercooled water and snow melt
- ◆ Ice sedimentation and time-split fall terms

# mp\_physics=14

## WDM 5-class scheme

- ◆ Version of WSM5 that is double-moment for warm rain processes
- ◆ 5-class microphysics with ice
- ◆ CCN, and number concentrations of cloud and rain also predicted

# mp\_physics=5

Ferrier (current NAM) scheme

- ◆ Designed for efficiency

- Advection only of total condensate and vapor
- Diagnostic cloud water, rain, & ice (cloud ice, snow/graupel) from storage arrays – assumes fractions of water & ice within the column are fixed during advection

- ◆ Supercooled liquid water & ice melt

- ◆ Variable density for precipitation ice (snow/graupel/sleet) – “rime factor”

# mp\_physics=6

## WSM 6-class scheme

- ◆ From Hong and Lim (2006, JKMS)
- ◆ 6-class microphysics with graupel
- ◆ Ice number concentration as in WSM3 and WSM5
- ◆ New combined snow/graupel fall speed
- ◆ Time-split fall terms with melting

# mp\_physics=16

## WDM 6-class scheme

- ◆ Version of WSM6 that is double-moment for warm rain processes
- ◆ 6-class microphysics with graupel
- ◆ CCN, and number concentrations of cloud and rain also predicted

# mp\_physics=7

Goddard 6-class scheme

- ◆ From Tao et al.
- ◆ 6-class microphysics with graupel
- ◆ Based on Lin et al. with modifications for ice/water saturation
- ◆ gsfcgce\_hail switch for hail/graupel properties
- ◆ gsfcgce\_2ice switch for removing graupel or snow processes
- ◆ Time-split fall terms with melting



# mp\_physics=8

New Thompson et al. scheme in V3.1

- ◆ Replacement of Thompson et al. (2007) scheme that was option 8 in V3.0
- ◆ 6-class microphysics with graupel
- ◆ Ice and rain number concentrations also predicted (double-moment ice)
- ◆ Time-split fall terms

# mp\_physics=98

Old Thompson et al. 2007 graupel scheme

- ◆ From Thompson et al. (2007)
- ◆ Was option 8 in Version 3.0
- ◆ 6-class microphysics with graupel
- ◆ Ice number concentration also predicted (double-moment ice)
- ◆ Time-split fall terms

# mp\_physics=10

Morrison 2-moment scheme

- ◆ New in Version 3.0
- ◆ 6-class microphysics with graupel
- ◆ Number concentrations also predicted for ice, snow, rain, and graupel (double-moment)
- ◆ Time-split fall terms

# no\_mp\_heating=1

- ◆ Turn off heating effect of microphysics
  - Zeroes out the temperature tendency
  - Equivalent to no latent heat
  - Other microphysics processes not affected
  - New in Version 3

# mp\_zero\_out

Microphysics switch (also mp\_zero\_out\_thresh)

- ◆ 1: all values less than threshold set to zero (except vapor)
- ◆ 2: as 1 but vapor also limited  $\geq 0$
- ◆ Note: this option will not conserve total water
- ◆ Not needed when using positive definite advection
- ◆ NMM: Recommend mp\_zero\_out=0

# nphs

- ◆ Time steps between microphysics calls
- ◆ Same as parameter for turbulence/PBL/LSM
- ◆ Typical value is chosen to give a frequency of 1-3 minutes, i.e.  $60/dt$  to  $180/dt$

# Microphysics Options

Recommendations about choice

- ◆ Probably not necessary to use a graupel scheme for  $dx > 10$  km
  - Updrafts producing graupel not resolved
  - Cheaper scheme may give similar results
- ◆ When resolving individual updrafts, graupel scheme should be used
- ◆ All domains use same option

# Rainfall Output

- ◆ Cumulus and microphysics can be run at the same time
- ◆ ARW outputs rainfall accumulations since simulation start time (0 hr) in mm
- ◆ RAINC comes from cumulus scheme
- ◆ RAINNC comes from microphysics scheme
- ◆ Total is RAINC+RAINNC
  - RAINNCV is time-step value
  - SNOWNC/SNOWNCV are snow sub-set of RAINC/RAINNCV (also GRAUPELNC, etc.)




# Rainfall Output

- ◆ Cumulus and microphysics can be run at the same time
- ◆ NMM outputs rainfall accumulations in mm
- ◆ TPREC controls zeroing out frequency
- ◆ ACPREC is the total precipitation
- ◆ CUPREC is the part that comes from the cumulus scheme
- ◆ The microphysics part is  $ACPREC - CUPREC$



# Physics Interactions

# Direct Interactions of Parameterizations



# Solver Calling Sequence (ARW example)

Call to solver advances one domain by one model time-step

- ◆ Physics tendencies

- Radiation, surface, land-state update, PBL, cumulus, grid-fdda, obs-fdda

- ◆ Dynamics tendencies

- Diffusion, advection, dynamics terms (for 3d momentum, theta, geopotential, surface pressure)

- ◆ Acoustic steps

- Update 3d momentum, theta, surface pressure, height

- ◆ Scalar dynamics tendencies and update

- Advection, diffusion of moist (qv,qc, etc.), scalar, tracer, tke, (and chemistry) variables

- ◆ Microphysics update

# &physics

Seven major physics categories:

`mp_physics: 0,1,2,3,4,5,6,8,10`

`ra_lw_physics: 0,1,3,99`

`ra_sw_physics: 0,1,2,3,99`

`sf_sfclay_physics: 0,1,2`

`sf_surface_physics: 0,1,2,3,99` (set before  
running `real` or `ideal`, need to match with  
`num_soil_layers` variable)

`ucm_call = 0,1`

`bl_pbl_physics: 0,1,2,99`

`cu_physics: 0,1,2,3,99`

# PBL schemes in V3.1

bl_pbl_physics	Scheme	Reference	Added
1	YSU	Hong, Noh and Dudhia (2006, MWR)	2004
2	MYJ	Janjic (1994, MWR)	2000
3	GFS	Hong and Pan (1996, MWR)	2005
4	QNSE	Sukoriansky, Galperin and Perov (2005, BLM)	2009
5	MYNN2	Nakanishi and Niino (2006, BLM)	2009
6	MYNN3	Nakanishi and Niino (2006, BLM)	2009
7	ACM2	Pleim (2007, JAMC)	2008
8	BouLac	Bougeault and Lacarrere (1989, MWR)	2009
99	MRF	Hong and Pan (1996, MWR)	2000

# PBL schemes in V3.1

3.2 changes

bl_pbl_physics	Scheme	Cores	sf_sfclay_physics	Prognostic variables	Diagnostic variables	Cloud mixing
1	YSU	ARW NMM	1		exch_h	QC,QI
2	MYJ	ARW NMM	2	TKE_MYJ	EL_MYJ, exch_h	QC,QI
3	GFS(hwrf)	NMM	3			QC,QI
4	QNSE	ARW NMM	4	TKE_MYJ	EL_MYJ, exch_h, exch_m	QC,QI
5	MYNN2	ARW	1,2,5	QKE	Tsq, Qsq, Cov, exch_h, exch_m	QC
6	MYNN3	ARW	1,2,5	QKE, Tsq, Qsq, Cov	exch_h, exch_m	QC
7	ACM2	ARW	1,7			QC,QI
8	BouLac	ARW	1,2	TKE_PBL	EL_PBL, exch_h, exch_m, wu_tur, wv_tur, wt_tur, wq_tur	QC
99	MRF	ARW NMM	1			QC,QI

# LES schemes in V3.1

Unified horizontal and vertical mixing (for  $dx \sim dz$ ).

Typically needed for  $dx < \sim 200$  m. Also use `mix_isotropic=1`.

bl_pbl_p physics	diff_opt	km_opt	Scheme	Cores	sf_sfclay _physics	isfflx	Prognostic variables
0	2	2	tke	ARW	0,1,2	0,1,2	tke
0	2	3	3d Smagorinsky	ARW	0,1,2	0,1,2	

## Namelist isfflx controls surface flux methods

isfflx	sf_sfclay_physics	Heat flux	Drag	Real/Ideal
0	0	From namelist tke_heat_flux	From namelist tke_drag_coefficient	Ideal
1	1,2	From LSM/sfclay physics (HFX, QFX)	From sfclay physics (UST)	Real
2	1,2	From namelist tke_heat_flux	From sfclay physics (UST)	Ideal



# Microphysics schemes in V3.2

mp_physics	Scheme	Reference	Added
1	Kessler	Kessler (1969)	2000
2	Lin (Purdue)	Lin, Farley and Orville (1983, JCAM)	2000
3	WSM3	Hong, Dudhia and Chen (2004, MWR)	2004
4	WSM5	Hong, Dudhia and Chen (2004, MWR)	2004
5	Eta (Ferrier)	Rogers, Black, Ferrier, Lin, Parrish and DiMego (2001, web doc)	2000
6	WSM6	Hong and Lim (2006, JKMS)	2004
7	Goddard	Tao, Simpson and McCumber (1989, MWR)	2008
8 (+98)	Thompson (+old)	Thompson, Field, Rasmussen and Hall (2008, MWR)	2009
9	Milbrandt 2-mom	Milbrandt and Yau (2005, JAS)	2010
10	Morrison 2-mom	Hong and Pan (1996, MWR)	2008
14	WDM5	Lim and Hong (2010,...)	2009
16	WDM6	Lim and Hong (2010,...)	2009

# Microphysics schemes in V3.2

mp_physics	Scheme	Cores	Mass Variables	Number Variables
1	Kessler	ARW	Qc Qr	
2	Lin (Purdue)	ARW	Qc Qr Qi Qs Qg	
3	WSM3	ARW	Qc Qr	
4	WSM5	ARW NMM	Qc Qr Qi Qs	
5	Eta (Ferrier)	ARW NMM	Qc Qr Qs (Qt*)	
6	WSM6	ARW NMM	Qc Qr Qi Qs Qg	
7	Goddard	ARW	Qc Qr Qi Qs Qg	
8 (/98)	Thompson(/old)	ARW NMM	Qc Qr Qi Qs Qg	Ni Nr (/Ni)
9	Milbrandt 2-mom	ARW	Qc Qr Qi Qs Qg Qh	Nc Nr Ni Ns Ng Nh
10	Morrison 2-mom	ARW	Qc Qr Qi Qs Qg	Nr Ni Ns Ng
14	WDM5	ARW	Qc Qr Qi Qs	Nn** Nc Nr
16	WDM6	ARW	Qc Qr Qi Qs Qg	Nn** Nc Nr

\* Advects only total condensate \*\* Nn= CCN number