



WRF Modeling System Overview

Jimmy Dudhia

What is WRF?

- WRF: Weather Research and Forecasting Model
 - Used for both research and operational forecasting
- It is a supported “community model”, i.e. a free and shared resource with distributed development and centralized support
- Its development is led by NCAR, NOAA/ESRL and NOAA/NCEP/EMC with partnerships at AFWA, FAA, and collaborations with universities and other government agencies in the US and overseas

What are ARW and NMM?

- The Advanced Research WRF (ARW) and Nonhydrostatic Mesoscale Model (NMM) are dynamical cores
 - Dynamical core includes mostly advection, pressure-gradients, Coriolis, buoyancy, filters, diffusion, and time-stepping
- Both are Eulerian mass dynamical cores with terrain-following vertical coordinates
- ARW support and development are centered at NCAR/MMM
- NMM development is centered at NCEP/EMC and support is provided by NCAR/DTC
- This tutorial is for both dynamical cores
- Both are downloadable in the same WRF tar file
- Physics, the software framework, and parts of data pre- and post-processing are shared between the dynamical cores

WRF as a Community Model

- Version 1.0 WRF was released December 2000
- Version 2.0: May 2004 (NMM added, EM nesting released)
- Version 2.1: August 2005 (EM becomes ARW)
- Version 2.2: December 2006 (WPS released)
- Version 3.0: April 2008 (includes global ARW version)
- Version 3.1: April 2009
- Version 3.2: April 2010
 - Version 3.2.1 August 2010
- Version 3.3 April 2011


What can WRF be used for?

- ARW and NMM
 - Atmospheric physics/parameterization research
 - Case-study research
 - Real-time NWP and forecast system research
 - Data assimilation research
 - Teaching dynamics and NWP
- ARW only
 - Regional climate and seasonal time-scale research
 - Coupled-chemistry applications
 - Global simulations
 - Idealized simulations at many scales (e.g. convection, baroclinic waves, large eddy simulations)

Who uses WRF?

- Academic atmospheric scientists (dynamics, physics, weather, climate research)
- Forecast teams at operational centers
- Applications scientists (e.g. Air Quality, Hydrology, Utilities)


WRF Modeling System Flow Chart



Modeling System Components

- WRF Pre-processing System (WPS)
 - Real-data interpolation for NWP runs
 - New obsgrid program for adding more obs to analysis
- WRF Model (ARW and NMM dynamical cores)
 - Initialization programs for real and (for ARW) idealized data (real.exe/ideal.exe)
 - Numerical integration program (wrf.exe)
- Graphics and verification tools including MET
- WRFDA (separate tutorial)
- WRF-Chem (separate tutorial)
- WRF-Fire – wildland model for forest fires

WPS and WRF Program Flow



Real-Data Applications

- Numerical weather prediction
- Meteorological case studies
- Regional climate
- Applications: air quality, wind energy, hydrology, etc.



Real-Data Applications

- Need time-independent information for chosen *domain* (simulation grid area)
- GEOGRID program
 - Map projection information
 - 2d gridded latitude, longitude, Coriolis parameter, map-scale factors, etc.
 - Topographic information
 - 2d gridded elevation, vegetation and soil categories, etc.



Real-Data Applications

- Need initial conditions (initial analysis time)
- UNGRIB and METGRID programs
 - 3d fields of horizontal wind, temperature, geopotential height, relative humidity
 - 2d fields of surface or sea-level pressure, surface temperature, relative humidity, horizontal winds
 - Time-dependent land-surface fields snow-cover, soil temperature, soil moisture



Real-Data Applications

- Regional domains need *specified* lateral boundary conditions at later times (e.g. every 6 hours) through forecast period
 - 3d fields of horizontal wind, temperature, geopotential height, water vapor
 - 2d field of surface pressure
- Long simulations (> 1 week) also need lower boundary condition at later analysis times
 - 2d fields of sea-surface temperature, sea-ice



Real-Data Applications

- Lateral Boundary Conditions (linear in time)
 - The *wrfbdy* file contains later gridded information at model points in a zone (e.g.) 5 points wide around the domain
 - The boundary fields are linearly time-interpolated from boundary times to the current model time
 - This specifies the outer values, and is used to nudge the next 4 interior points
- Lower Boundary Condition (step-wise)
 - New SSTs are read in and overwritten at each analysis time from *wrflowinp* file



Real-Data Applications


Summary

- Pre-processing for regional domains therefore needs multiple times for lateral boundary conditions during whole forecast period (UNGRIB and METGRID should be run for all needed analysis times)
 - Note: Global models only need initial analysis
 - Real-time regional NWP often uses global forecast for boundary conditions
- Long simulations also need lower boundary information on SST and sea ice to update them over periods of weeks, months, years



Nesting

- Running multiple domains with increasing resolution in nested areas
- Parent has *specified* boundary conditions from *wrfbdy* file
- *Nested* boundary conditions come from parent



Nesting (Two-Way)

- Lateral boundary condition is provided by parent domain at every parent step
- Method is same as for outer domain (specified and relaxation zones)
- Additional fields include vertical motion and microphysics species
- Feedback: Interior of nest overwrites overlapped parent area



Nesting (Two-Way)

- Sequence
 - Parent domain runs a time-step to $t+dt$
 - Nest boundaries from beginning and end of time-step interpolated
 - Nest runs typically three steps ($dt/3$) using time-interpolated parent info at nest boundaries
 - After nest reaches $t+dt$, feedback overwrites parent in overlapped region
 - Repeat




One-Way Nesting

- As two-way nesting but no feedback
- Can also be done with NDOWN program to take a previous WRF run output and provide nest boundary conditions at parent output frequency
 - Uses parent WRF run instead of analysis for initial and lateral boundary conditions



WPS and WRF Program Flow



WPS Functions

- Define simulation domain area (and nests)
- Produce terrain, landuse, soil type etc. on the simulation domain (“static” fields)
- De-grib GRIB files for meteorological data (u, v, T, q, surface pressure, soil data, snow data, sea-surface temperature, etc.)
- Interpolate meteorological data to WRF model grid (horizontally)
- Optionally add more observations to analysis (separate obsgrid program)

WRFDA (Data Assimilation)

- Variational data assimilation (3D-Var and 4D-Var)
- Ensemble DA
- Hybrid variational/ensemble DA

Function

- Ingest observations to improve WRF input analysis from WPS
- May be used in cycling mode for updating WRF initial conditions after WRF run
- Also used for observation impact data studies

WRFDA

- Supported for data types
 - Conventional surface and upper air, wind profiler, aircraft
 - Remote sensing data: Cloud-tracked winds, satellite-retrieved profiles, ground-based/satellite GPS, scatterometer ocean surface winds, radar radial velocity and reflectivity, satellite radiance
- Background error covariance for ARW from
 - NMC method
 - Ensemble method

WRF-Chem

- Supported by NOAA/ESRL
- Includes chemistry species and processes, many chemistry options
- Also needs emissions data
- Included in WRF tar file, but requires special compilation option

WRF real and ideal functions

- REAL
 - Creates initial and boundary condition files for real-data cases
 - Does vertical interpolation to model levels (when using WPS)
 - Does vertical dynamic (hydrostatic) balance
 - Does soil vertical interpolations and land-use mask checks
- IDEAL (ARW only)
 - Programs for setting up idealized case
 - Simple physics and usually single sounding
 - Initial conditions and dynamic balance


WRF Model

- WRF
 - Dynamical core (ARW or NMM) is compile-time selectable
 - Uses initial conditions from REAL or IDEAL (ARW)
 - Real-data cases use boundary conditions from REAL
 - Runs the model simulation with run-time selected namelist switches (such as physics choices, timestep, length of simulation, etc.)
 - Outputs history and restart files

Dynamical Cores

- Dynamical cores include
 - Advection, Coriolis, pressure gradient terms, diffusion, in the basic equations
 - Finite differencing for these terms on a grid and time-stepping method and numerical filters
- ARW dynamical core
 - Designed for research as well as NWP
 - Idealized cases, regional climate, WRF-Chem, more options
- NMM dynamical core
 - Used in NCEP operational regional models (i.e. NAM and HWRF)
 - Focused on NWP applications, more limited options

WRF Modeling System Flow Chart





Basic Software Requirement

- Fortran 90/95 compiler
 - Code uses standard f90 (very portable)
- C compiler
 - “Registry”-based automatic Fortran code generation (for argument lists, declarations, nesting functions, I/O routines)
- Perl
 - configure/compile scripts
- netcdf library
 - for I/O (other I/O formats semi-supported)
- Public domain mpich for MPI
 - if using distributed memory option


User Support

- Email: wrfhelp@ucar.edu
- User Web pages:
 - ARW: <http://www.mmm.ucar.edu/wrf/users/>
 - NMM: <http://www.dtcenter.org/wrf-nmm/users/>
 - Latest update for the modeling system
 - WRF software download
 - Various documentation
 - Users' Guide
 - Technical Note (ARW Description)

ARW Hurricane Katrina Simulation (4km)



ARW Convective-scale Forecasting (4km)



Tutorial Schedule

- Lectures for WRF: Mon.-Fri.
- Practice for WRF: Mon.-Fri.
 - 2 Groups (a.m./p.m.)
- WRF-Chem tutorial: Next Mon.-Tue.
- WRF-DA tutorial: Next Wed.-Fri.