

Introduction to WRF-Chem

Georg Grell

Steven E. Peckham, Stuart A. McKeen + others from NOAA/ESRL

Jerome Fast, William Gustafson jr., + many others from PNNL

+ Saulo Freitas, Karla Longo (CPTEC, BRAZIL)

+ Christine Wiedinmyer, Xue-Xi, Gabi Pfister, Mary Barth and many others from
NCAR

**+ many more national and international
collaborators**

WRF-Chem web site - <http://wrf-model.org/WG11>





WRF-Chem

Community effort

**Largest contributing groups: ESRL,
PNNL, NCAR**

**Other significant contributions
from: University of Chile, CPTEC
Brazil, University of Fairbanks,
NASA**



WRF-Chem

The help desk:

wrfchemhelp.gsd@noaa.gov




WRF-Chem

- Online, completely embedded within WRF CI
- Consistent: all transport done by meteorological model
 - Same vertical and horizontal coordinates (no horizontal and vertical interpolation)
 - Same physics parameterization for subgrid scale transport
 - No interpolation in time
- Easy handling (Data management)
- Ideally suited to study feedbacks between chemistry and meteorology
- Ideally suited for air quality forecasting on regional to cloud resolving scales

Why Online?

- In models, with increasing horizontal resolution, the variability of the vertical velocity becomes much more important
- Offline might introduce a large error in estimate of vertical mass transport



Gas Phase Chemistry Packages


- Chemical mechanism from RADM2 (Quasi Steady State Approximation method with 22 diagnosed, 3 constant, and 38 predicted species is used for the numerical solution)
- Carbon Bond (CBM-Z) based chemical mechanism, and the
- Kinetic PreProcessor (KPP) – Many different equations files exist
 - RADM2, versions of RACM, MOZART, CBMZ, SAPRC99,...

Available Aerosols modules

1. PM advection, transport, emissions and deposition only
2. GOCART
3. Modal approach (MADE/SORGAM)
4. Sectional approach (MOSAIC)


Aerosol modules comparison

(1) Modal



composition
sulfate
nitrate
ammonium
chloride
carbonate
sodium
calcium
other inorganics
organic carbon
elemental carbon

(2) Sectional



(3) GOCART: Sections for dust and sea salt, otherwise total mass only



For NWP a bulk scheme is very attractive

GOCART (Currently used in real-time FIM-Chem, RR-Chem, and HRRR-Chem)

- Much simpler than the sectional and model schemes
 - Calculates only with the total mass of the aerosol components
 - Provides no information on
 - Particle size
 - Particle concentration
 - E.g., when particles grow, the aerosol mass increases but we don't know how their size/number changes
- Numerically very efficient
- Coupled with radiation (Mie scattering and extinction calculations)



For research on aerosol direct and indirect effects modal and sectional approaches are more attractive

Less assumptions are made when coupled to atmospheric radiation and/or microphysics



Selection of radiation parameterizations for aerosol direct effect

For V3.3.1 all aerosol modules were hooked up to Goddard short wave radiation, and RRTMG short and long wave scheme.

More to come for V3.4



Selection of microphysics parameterizations for aerosol indirect effect

For V3.3.1

Modal and sectional scheme only
can be used in combination with a
version of the Lin et al. Microphysics
scheme as well as the Morrison
scheme



Photolysis Packages – all coupled to aerosols and hydrometeors

- Madronich Photolysis
- Madronich F-TUV
- Fast-j photolysis scheme

Biogenic emissions

- May be calculated “online” based on USGS landuse
 - Easy to use
- May be input
- BEISv3.13 (offline reference fields, online modified)
 - Good choice, but difficult to use
- Use of MEGAN
 - Best choice!!



Model of Emissions of Gases and Aerosols from Nature (MEGAN) in WRF-Chem

Global, high resolution biogenic emissions


**Out of available biogenic emissions
modules only BEIS and MEGAN are actively
being worked on (developed)**

Preprocessor for MEGAN exists and can be downloaded
from NCAR

Fire Plumerise - Collaboration with Saulo Freitas from CPTEC in Brazil

1-D Cloud model to determine injection height


Satellite information (other aerial and ground observations may also be used) to determine fire location and fire properties




New in V3.3: Coupling of Aerosols to RRTMG Radiation (implemented by PNNL)

- Extended modular optical property module to compute information needed for both shortwave and longwave RRTMG radiation scheme
- Works for all aerosol modules
- Evaluated using AOD and extinction profile data over northern Africa associated with Saharan dust
 - GOCART dust emission module also extended to work with MADE/SORGAM and MOSAIC
 - See *Zhao et al., ACP, 2010* for more details

Dust Emissions from 2 Treatments




AOD under Various Scenarios – Dust Emissions and Aerosol Models





Aerosol-Cloud Aerosol Interactions: What's new in V3.3

- Morrison microphysics scheme now coupled with either prescribed aerosols or predicted aerosols from both MADE/SORGAM or MOSAIC (PNNL + Hugh Morrison(NCAR))
- Using same treatment for aerosol activation as implemented previously with Lin microphysics scheme, as described in *Chapman et al., ACP, 2009*
- Current applications (publications in progress):
 - Effect of aerosols on marine stratocumulus during VOCALS-Rex



➤ Effect of aerosols on mixed phase clouds during ISDAC / ARCTAS



droplet number produced by aerosol activation scheme coupled with Morrison microphysics consistent with aircraft data


Secondary Organic Aerosols: New in V3.3

Added 'volatility basis set' approach to MOSAIC (PNNL)

- Based on *Robinson et al. (2007)* and *Shrivastava et al. (2008)*
- Coupled with SAPRC99 photochemistry – which has also been added as a separate stand alone package, in collaboration with Pablo Saide (Univ. Iowa)
- Release version uses 2 volatility bins for simplicity and 4 size bins, 8 volatility bins approach available on request (computationally expensive)
- Oxygen and carbon species treated separately to compute O:C ratios
- Anthropogenic, biomass burning, and biogenic sources of organics tracked

Like all SOA parameterizations, it should be treated with caution


- Predictions not perfect, but better than simulating non-volatile POA only
- Evaluated against surface and aircraft data collected during MILAGRO field campaign – see *Shrivastava et al., ACPD, 2010* and *Fast et al. ACP 2009* for more details
- Optical properties for SOA treated the same as POA – for now
- Aerosol-cloud interactions not treated for SOA presently



DMS and Sea-Salt Emissions

- DMS chemistry now included in GOCART and MOSAIC

SO₂ over the Southeastern Pacific Ocean during VOCALS-Rex, looking Southeast



Volcanic ash in WRF-Chem V3.3

Collaboration with University of Alaska in
Fairbanks as well as INPE/CPTEC in Brazil,
Publications in progress

Impact of Volcanoes

- Ash-fall near eruption
- Transport of fine ash in high concentrations for long distances
- Impact on weather, climate, and air quality



The plume of the 30 Sept/1 Oct 1994 eruption of Kliuchevskoi Volcano, Kamchatka taken from the space shuttle STS-68 mission (Russia)

ASH Volcanoes Prediction

Based on Mastin et al. (2009) dataset

1. 1535 volcanoes with lat, lon, elevation, eruption classification (ESP)
2. Table describing injection height, duration, eruption rate, volume and mass fraction (<63um)

| ESP | Type | Example | H km above vent | Duration hr | Eruption rate (kg/s) | Volume (km3) | mass fraction less than 63 micron |
|-----|-----------------------|---|-----------------|-------------|----------------------|--------------|-----------------------------------|
| | | Cerro Negro, Nicaragua, 4/13/1992 | 7 | 60 | 1,E+05 | 0,01 | 0,05 |
| M0 | Standard mafic | Etna, Italy, 7/19-24/2001 | 2 | 100 | 5,E+03 | 0,001 | 0,02 |
| M1 | small mafic | Cerro Negro, Nicaragua, 4/9-13/1992 | 7 | 60 | 1,E+05 | 0,01 | 0,05 |
| M2 | medium mafic | Fuego, Guatemala, 10/14/1974 | 10 | 5 | 1,E+06 | 0,17 | 0,1 |
| M3 | large mafic | Spurr, USA, 8/18/1992 | 11 | 3 | 4,E+06 | 0,015 | 0,4 |
| S0 | standard silicic | Ruapehu, New Zealand, 6/17/1996 | 5 | 12 | 2,E+05 | 0,003 | 0,1 |
| S1 | small silicic | Spurr, USA, 8/18/1992 | 11 | 3 | 4,E+06 | 0,015 | 0,4 |
| S2 | medium silicic | St. Helens, USA, 5/18/1980 | 15 | 8 | 1,E+07 | 0,15 | 0,5 |
| S3 | large silicic | St. Helens, USA, 5/18/1980 (pre-9 AM) | 25 | 0,5 | 1,E+08 | 0,05 | 0,5 |
| S8 | co-ignimbrite silicic | Soufrière Hills, Montserrat (composite) | 10 | 0,01 | 3,E+06 | 0,0003 | 0,6 |
| S9 | Brief silicic | none | 0 | -- | -- | -- | -- |
| U0 | default submarine | | | | | | |


10 size bins for prediction of **ash-fall** and transport of volcanic ash

| Particle Size Bin | Phi | Percentage of mass |
|-------------------------------|--------|--------------------|
| 1 – 2mm | -1 – 0 | 2 |
| 0.5 – 1 mm | 0 – 1 | 4 |
| 0.25 – 0.5 mm | 1 – 2 | 11 |
| 125 – 250 μm | 2 – 3 | 9 |
| 62.5 – 125 μm | 3 – 4 | 9 |
| 31.25 – 62.5 μm | 4 – 5 | 13 |
| 15.625 – 31.25 μm | 5 – 6 | 16 |
| 7.8125 – 15.625 μm | 6 – 7 | 16 |
| 3.9065 – 7.8125 μm | 7 – 8 | 10 |
| < 3.9 μm | > 8 | 10 |


4 size bins for prediction if transport only is of interest

| Particle Size Bin | Phi | Percentage of mass |
|-------------------------------|-------|--------------------|
| 15.625 – 31.25 μm | 5 – 6 | 16 |
| 7.8125 – 15.625 μm | 6 – 7 | 16 |
| 3.9065 – 7.8125 μm | 7 – 8 | 10 |
| < 3.9 μm | > 8 | 10 |

Tephra-fall deposits (g/m^2)
Redoubt Volcano, south-central Alaska
December 15, 1989



Observed



VOLCANIC ASH FALL (g/m^2)




0 8 16 24 32 40 48 56 64 72 80 88 96

Predicted by WRF-Chem


First WRF-Chem runs for “Big E”

- 30km horizontal resolution
- 10 ash bins
- Ash settling, dry deposition, and wet deposition included
- Aerosol optical properties easily implemented for ash


Comparison of ash forecasts (London VAAC and WRF-Chem) at 0000Z, April 15



VA advisory
from London




Forecast compared to Munich Lidar, April 17, 06Z



A

Munich at 48.148 Lat 11.573 long, total ash April 17

lid: 2010 04 14_00:00:00




Ash Concentration ($\mu\text{g}/\text{m}^3$)

| | | | | | | |
|----|----|-----|-----|-----|-----|-----|
| 20 | 60 | 100 | 140 | 180 | 220 | 260 |
|----|----|-----|-----|-----|-----|-----|

B

2010-04-17_06:00:00 48.148 Latitude 11.573 Longitude



C





How is the meteorological forecast affected by aerosol?

- In general large importance for climate simulations is recognized (when integrating models over 100's of years, small differences in the earth's energy budget are extremely important)
- Weather forecasting for only a few days?
 - Much research needed, but direct effect may positively influence forecasts when strong signals exist
 - Influence on meteorological data assimilation

No fires

24-hr forecast

With fires







Observed (black) and predicted (blue) sounding for Fairbanks, Alaska, on July 4, 0000UTC.

Chemical data assimilation: WRF-Chem and PM2.5

2 months worth of WRF-Chem runs:

1. New England 2004 to estimate background error covariances and lengthscales
2. Houston 2006 for evaluation





Chemical data assimilation: ARW-WRF/Chem and PM2.5

Much work in progress at ESRL (EnKF) as well as at NCAR (AOD assimilation with GSI), not released to community yet

If you need chemical data assimilation, contacts: wrfchemhelp desk will tell you who and when

WRF-Chem current and future work – ESRL/ GSD, CSD – V3.4

- Simple aqueous chemistry and wet deposition for resolved precipitation as well as convective parameterization (CMAQ module)
- CO₂ emissions module- Includes a high spatiotemporal resolution biospheric flux model –Vegetation and Photosynthesis (uses MODIS reflectances as input)
- SOA: the volatility basis set approach has been coupled with modal aerosol scheme (additional option: MADE/VBSSOA)



WRF/Chem current and future work – ESRL + other groups – V3.4, V3.4.x

- CH₄ emissions module
 - Different CH₄ tracers (anthropogenic, biospheric, ...)
 - Several CH₄ flux models are implemented: Wetland fluxes (Kaplan, 2002), Soil uptake (Ridgwell et al., 1999), Termite fluxes (Sanderson, 1996)
- Dust parameterization from AFWA (S. Jones and G Creighton) is working and is being evaluated
- Volcanic SO₂ emissions
- Coupling of volcanic emissions with other aerosol modules
- Aerosols will be coupled with convective parameterization (G₃, collaboratively with S. Freitas)
- Using WPS to run WRF-Chem off global FIM-Chem

WRF/Chem current and future work – NCAR/ACD

– V3.4

- Add wet scavenging of gases (see talk by Gabi Pfister, Thursday, June 23)
- Improve SOA gas chemistry and add SOA aqueous chemistry (likely to be hooked with MOZCART)
- **Upper Chemical Boundary Conditions**
Chemical UBC are taken from WACCM climatology for past, present and future (talk by M. Barth et al., Thursday, June 23)
- **Reduced Chemistry**
Implemented reduced chemical mechanism (Howeling et al., 1998); useful for long climate runs and compatible with CAM-Chem (-> talk by Jerome Fast in Physics Section, Thursday, June 23)
- **Aircraft Tracking Tool**
Enables output for specified location in model time steps

WRF-Chem ongoing and future work – PNNL – V3.4.x , V3.5 or later

- Interactive physics: More microphysics, radiation schemes, also a new inexpensive modal approach
- Continued work on cloud-aerosol interactions
- new aerosol model is planned (MOSAIC-ext), that simulates the evolution of the transition between internal and external aerosol mixing states
- ice-aerosol interactions to be included

WRF-Chem ongoing and future work – PNNL

- Aerosol modeling test bed is making progress

<http://www.pnl.gov/atmospheric/research/aci/amt/index.stm>

- Some of the Analysis Toolkit Software available via the web site
- MILAGRO test bed data is finished,
- CHAPS, VOCALS, ISDAC/ARCTAS, CARES/CalNex integrated datasets (field campaign + routine monitoring) planned for the future

Resources

- WRF project home page
 - <http://www.wrf-model.org>
- WRF users page (linked from above)
 - <http://www.mmm.ucar.edu/wrf/users>
- On line documentation (also from above)
 - http://www.mmm.ucar.edu/wrf/WG2/software_v2
- WRF users help desk
 - wrfhelp@ucar.edu
- WRF-Chem users help desk
 - wrfchemhelp.gsd@noaa.gov