



# WRF Modeling System Overview


Jimmy Dudhia



# What is WRF?

---

- WRF: Weather Research and Forecasting Model
  - Used for both research and operational forecasting
- It is a supported “community model”, i.e. a free and shared resource with distributed development and centralized support
- Its development is led by NCAR, NOAA/ESRL and NOAA/NCEP/EMC with partnerships at AFWA, FAA, DOE/PNNL and collaborations with universities and other government agencies in the US and overseas



# What is ARW?

---


- WRF has two dynamical cores: The Advanced Research WRF (ARW) and Nonhydrostatic Mesoscale Model (NMM)
  - Dynamical core includes mostly advection, pressure-gradients, Coriolis, buoyancy, filters, diffusion, and time-stepping
- Both are Eulerian mass dynamical cores with terrain-following vertical coordinates
- ARW support and development are centered at NCAR/MMM
- NMM development is centered at NCEP/EMC and support is provided by NCAR/DTC
- This tutorial is for only the ARW core
- Both are downloadable in the same WRF tar file
- Physics, the software framework, and parts of data pre- and post-processing are shared between the dynamical cores



# WRF as a Community Model

---


- Version 1.0 WRF was released December 2000
- Version 2.0: May 2004 (NMM added, EM nesting)
- Version 2.1: August 2005 (EM becomes ARW)
- Version 2.2: December 2006 (WPS released)
- Version 3.0: April 2008 (includes global ARW version)
- Version 3.1: April 2009
- Version 3.2: April 2010
- Version 3.3: April 2011
- Version 3.4: April 2012
- Version 3.5: April 2013
  - Version 3.5.1 September 2013 (current version)



# What can WRF be used for?

---

- ARW and NMM
  - Atmospheric physics/parameterization research
  - Case-study research
  - Real-time NWP and forecast system research
  - Data assimilation research
  - Teaching dynamics and NWP
- ARW only
  - Regional climate and seasonal time-scale research
  - Coupled-chemistry applications
  - Global simulations
  - Idealized simulations at many scales (e.g. convection, baroclinic waves, large eddy simulations)




# Who uses WRF?

---

- Academic atmospheric scientists (dynamics, physics, weather, climate research)
- Forecast teams at operational centers
- Applications scientists (e.g. Air Quality, Hydrology, Utilities)



# WRF Modeling System Flow Chart



# Modeling System Components


---

- WRF Pre-processing System
  - Real-data interpolation for NWP runs (WPS)
  - Program for adding more observations to analysis (obsgrid)
- WRF Model (ARW and NMM dynamical cores)
  - Initialization programs for real and (for ARW) idealized data (real.exe/ideal.exe)
  - Numerical integration program (wrf.exe)
- Graphics and verification tools including MET
- WRFDA (separate tutorial)
- WRF-Chem (separate tutorial)
- WRF-Fire – wildland model for forest fires



# WPS and WRF Program Flow


---



# Real-Data Applications

---


- Numerical weather prediction
- Meteorological case studies
- Regional climate
- Applications: air quality, wind energy, hydrology, etc.



# Real-Data Applications

---

- Need time-independent information for chosen *domain* (simulation grid area)
- GEOGRID program
  - Map projection information
    - 2d gridded latitude, longitude, Coriolis parameter, map-scale factors, etc.
  - Topographic information
    - 2d gridded elevation, vegetation and soil categories, etc.



# Real-Data Applications

---


- Need time-dependent information
- Initial conditions (initial analysis time)
- Boundary conditions (later times)
  - except if running WRF globally
- UNGRIB and METGRID programs
  - 3d fields of horizontal wind, temperature, geopotential height, relative humidity
  - 2d fields of surface or sea-level pressure, surface temperature, relative humidity, horizontal winds
  - Time-sensitive land-surface fields: snow-cover, soil temperature, soil moisture



# Real-Data Applications

---


- Regional domains need *specified* lateral boundary conditions at later times (e.g. every 6 hours) through forecast period
  - 3d fields of horizontal wind, temperature, geopotential height, water vapor
  - 2d field of surface pressure
- Long simulations ( $> 1$  week) also need lower boundary condition at later times
  - 2d fields of sea-surface temperature, sea-ice, vegetation fraction



# Real-Data Applications

---

- Lateral Boundary Conditions (linear in time)
  - The *wrfbdy* file contains later gridded information at model points in a zone (e.g.) 5 points wide around the domain
  - The boundary fields are linearly time-interpolated from boundary times to the current model time
  - This specifies the outer values, and is used to nudge the next 4 interior points
- Lower Boundary Condition (step-wise)
  - New SSTs are read in and overwritten at each analysis time from *wrflowinp* file




# Real-Data Applications

---

## Summary



- Pre-processing for regional domains therefore needs multiple times for lateral boundary conditions during whole forecast period (UNGRIB and METGRID should be run for all needed analysis times)
  - Note: Global models only need initial analysis
  - Real-time regional NWP often uses global forecast for boundary conditions
- Long simulations also need lower boundary information on SST and sea ice to update them over periods of weeks, months, years



# Nesting

---

- Running multiple domains with increasing resolution in nested areas
- Parent has *specified* boundary conditions from *wrfbdy* file
- *Nested* boundary conditions come from parent



# Nesting (Two-Way)

---


- Lateral boundary condition is provided by parent domain at every parent step
- Method is same as for outer domain (specified and relaxation zones)
- Additional fields include vertical motion and microphysics species
- Feedback: Interior of nest overwrites overlapped parent area



# Nesting (Two-Way)

---


- Sequence
  - Parent domain runs a time-step to  $t+dt$
  - Nest boundaries from beginning and end of time-step interpolated
  - Nest runs typically three steps ( $dt/3$ ) using time-interpolated parent info at nest boundaries
  - After nest reaches  $t+dt$ , feedback overwrites parent in overlapped region
  - Repeat



# One-Way Nesting

---


- As two-way nesting but no feedback
- Can also be done with NDOWN program to take a previous WRF run output and provide nest boundary conditions at parent output frequency
  - Uses parent WRF run instead of analysis for initial and lateral boundary conditions



# WPS Functions

---


- Define simulation domain area (and nests)
- Produce terrain, landuse, soil type etc. on the simulation domain (“static” fields)
- De-grib GRIB files for meteorological data (u, v, T, q, surface pressure, soil data, snow data, sea-surface temperature, etc.)
- Interpolate meteorological data to WRF model grid (horizontally)
- Optionally add more observations to analysis (separate obsgrid program)




# WPS Data

---



- Geogrid: We provide elevation, land-use, soil type data (static fields)
  - Or user can input own static data in same easy-to-write format
- Metgrid: Supports input of time-dependent data (dynamic fields)
  - UNGRIB can provide these from GriB files
  - Or user can input own data in same “intermediate format” (simple binary files)



# WPS and WRF Program Flow



# Data Flow



# Dynamical Cores

---

- Dynamical cores include
  - Basic Dynamical Equations: Advection, Coriolis, pressure gradient terms, buoyancy, diffusion
  - Finite differencing: Staggered grid-structure, time-stepping method and numerical filters
- ARW dynamical core
  - Designed for research as well as NWP
  - Idealized cases, regional climate, WRF-Chem, more options
- NMM dynamical core
  - Used in NCEP operational regional models (i.e. NAM and HWRF)
  - Focused on NWP applications, more limited options



# WRFDA (Data Assimilation)

---

- Variational data assimilation (3D-Var and 4D-Var)
- Ensemble DA
- Hybrid variational/ensemble DA

## Function

- Ingest observations to improve WRF input analysis from WPS
- May be used in cycling mode for updating WRF initial conditions after WRF run
- Also used for observation impact data studies



# WRF-Chem

---

- Supported by NOAA/ESRL
- Includes chemistry species and processes, many chemistry options
- Also needs emissions data
- Included in WRF tar file, but requires special compilation option



# ARW Dynamics

## Key features:




- Fully compressible, non-hydrostatic (with hydrostatic option)
- Mass-based terrain following coordinate,  $\sigma$

$$\eta = \frac{(\pi - \pi_t)}{\mu}, \quad \mu = \pi_s - \pi_t$$

where  $\pi$  is hydrostatic pressure,

$\mu$  is column mass

- Arakawa C-grid staggering




# ARW Model

---

Key features:

- 3rd-order Runge-Kutta time integration scheme
- High-order advection scheme
- Scalar-conserving (positive definite option)
- Complete Coriolis, curvature and mapping terms
- Two-way and one-way nesting



# ARW Model

---

## Key features:


- Choices of lateral boundary conditions suitable for real-data and idealized simulations
  - Specified, Periodic, Open, Symmetric, Nested
- Full physics options to represent atmospheric radiation, surface and boundary layer, and cloud and precipitation processes
- Grid-nudging and obs-nudging (FDDA)
- Digital Filter Initialization option



# Graphics and Verification Tools

---

- ARW and NMM
  - RIP4 (Read, Interpolate and Plot)
  - Unified Post-Processor (UPP)
    - Conversion to GriB (for GrADS and GEMPAK)
  - MET (Model Evaluation Toolkit)
- ARW
  - NCAR Graphics Command Language (NCL)
  - ARWpost
    - Conversion program for GrADS
  - VAPOR (3D visualization tool)
  - IDV (3D visualization tool)



# Basic Software Requirement

---


- Fortran 90/95 compiler
  - Code uses standard f90 (very portable)
- C compiler
  - “Registry”-based automatic Fortran code generation (for argument lists, declarations, nesting functions, I/O routines)
- Perl
  - configure/compile scripts
- netcdf library
  - for I/O (other I/O formats semi-supported)
- Public domain mpich for MPI
  - if using distributed memory option



# Code Layers


---

- Top-level (framework): allocates space, handles nested domains and interpolation/feedback functions, time-stepping, solver calls, and i/o file contents and calls
- Intermediate level: “start” routine for initial calls, “solve” routine for run-time advancing, MPI handling
- Low-level: science code in plain Fortran (no MPI or I/O calls)





# WRF Hierarchical Software Architecture

- Driver Layer
  - Memory allocation, nest starting, time-stepping, I/O
- Mediation Layer
  - Solver
- Model Layer
  - Dynamics, physics



## WRF Two-Layer Domain Decomposition (patches, tiles, halo)


- **Single version of code enabled for efficient execution on:**
  - Shared-memory multiprocessors
  - Distributed-memory multiprocessors
  - Distributed clusters of SMPs
  - Vector and scalar processors



# Registry File

---

- Designed to make adding arrays or new namelist parameters easy
- Also can add them to “halo” for MPI communications (only sometimes needed)
- Allocates, passes, and declares, listed arrays for nesting, i/o and “solver” routines
  - Solver advances one domain by one time step
    - From solver, it can be passed to parts of the low-level code via argument lists



# User Support

---

- Email: wrfhelp@ucar.edu
- User Web pages:
  - ARW: <http://www.mmm.ucar.edu/wrf/users/>
  - NMM: <http://www.dtcenter.org/wrf-nmm/users/>
  - Latest update for the modeling system
  - WRF software download
  - Various documentation
    - Users' Guides (both cores)
    - Technical Note (ARW Description)
    - Technical Note (NMM Description)

