


Overview of Physical Parameterizations

Song-You Hong

(KIAPS: Korea Institute of Atmospheric Prediction Systems)

(Also, an NCAR affiliate scientist)

1) Concept



* Physical process in the atmosphere

- : Specification of heating, moistening and frictional terms in terms of dependent variables of prediction model
- Each process is a specialized branch of atmospheric sciences.

* Parameterization


The formulation of physical process in terms of the model variables as parameters, i.e., constants or functional relations.

1) Concept...continued

Subgrid scale process

Any numerical model of the atmosphere must use a finite resolution in representing continuum certain physical & dynamical phenomena that are smaller than computational grid.

- Subgrid process (Energy perspective)





$\Delta x \rightarrow 0$, the energy dissipation takes place by molecular viscosity
?

Objective of subgrid-scale parameterization

“To design the physical formulation of energy sink, withdrawing the equivalent amount of energy comparable to cascading energy down at the grid scale in an ideal situation.”


1) Concept...continued

* Parameterization that are only somewhat smaller than the smallest resolved scales.



where truncation limit ; spectral gap

Unfortunately, there is no spectral gap



2) Subgrid scale process & Reynolds averaging


Consider prognostic water vapor equation

$$\frac{\partial \rho q}{\partial t} = -\frac{\partial \rho u q}{\partial x} - \frac{\partial \rho v q}{\partial y} - \frac{\partial \rho w q}{\partial z} + \rho E - \rho C \quad \dots(1)$$

In the real atmosphere,

$$u = \bar{u} + u', \quad q = \bar{q} + q' \quad \left(\begin{array}{l} \text{*} \bar{a} : \text{grid-resolvable} \\ a' : \text{subgrid scale perturbation} \end{array} \right)$$

ρ' is neglected



2) Subgrid scale process & Reynolds averaging...continued

* Rule of Reynolds average : $\bar{q}' = 0, \bar{u}'\bar{q} = 0, \bar{u}\bar{q}' = \bar{u}\bar{q}$

then eq.(1) becomes

$$\frac{\partial \rho \bar{q}}{\partial t} = -\frac{\partial \rho \bar{u} \bar{q}}{\partial x} - \frac{\partial \rho \bar{v} \bar{q}}{\partial y} - \frac{\partial \rho \bar{w} \bar{q}}{\partial z} - \frac{\partial \bar{\rho} u' q'}{\partial x} - \frac{\partial \bar{\rho} v' q'}{\partial y} - \frac{\partial \bar{\rho} w' q'}{\partial z} + \rho E - \rho C \dots(2)$$

..... :

① ②

- ① grid-resolvable advection (dynamical process)
- ② turbulent transport

* How to parameterize the effect of turbulent transport

a) $-\rho \bar{w}' \bar{q}' = 0$: 0th order closure

b) $-\rho \bar{w}' \bar{q}' = K \frac{\partial \bar{q}}{\partial z}$: 1st order closure (K-theory)

c) obtain a prognostic equation for $\bar{w}' \bar{q}'$ from (1), (2)

$$\frac{\partial \rho w q}{\partial t} = -\frac{\partial \rho u w q}{\partial x} + \dots$$


taking Reynolds averaging,

$$\frac{\partial \overline{\rho w' q'}}{\partial t} = \frac{\partial \overline{\rho w' w' q'}}{\partial z}$$

$$-\rho \overline{w' w' q'} = K' \frac{\partial \overline{\rho w' q'}}{\partial z} \quad : \text{2nd order closure}$$


3) Schematic of physics algorithm : In modeled atmosphere : 6* ~9

$$\frac{d \ln \theta}{dt} = \frac{H}{c_p T}, \quad \frac{dq}{dt} = S, \quad \frac{d\vec{u}}{dt} = \nabla_z \vec{\tau}$$



1. Radiation

1.1 Concept





$$\text{TOA : } S = 1360 \text{ } Wm^{-2}$$

$$\text{Mean Flux : } \frac{S}{4} = 340 \text{ } Wm^{-2} \rightarrow \text{Energy source for Earth}$$


30% : reflected from the atmosphere clouds

→ Back to space by terrestrial infrared radiation

{
25% : absorbed in the atmosphere
45% : absorbed at the earth surface }



1.2 Solar radiative transfer



- At TOA,



$$F = S \left(\frac{dm}{d} \right)^2 \cos \theta_0 \quad (\theta_0 : \text{Zenith angle})$$

(Insolation)

- Basic equations $\mu = \cos \theta$



$$\mu \frac{dI(\tau, \mu, \phi)}{d\tau} = I(\tau, \mu, \phi) - J(\tau, \mu, \phi)$$

absorption source emission



$$d\tau = -k_v \rho_a dz \quad \tau(\text{optical depth}) = \int_z^{\infty} k_v(z') \rho_a(z') dz'$$

$$= \int_0^p k_v(p') q(p') \frac{dp'}{g}$$



$$J = J(\tau, \mu, \phi) = \frac{\tilde{\omega}}{4\pi} \int_0^{2\pi} \int_{-1}^1 I(\tau, \mu', \phi') P(\mu, \phi; \mu', \phi') d\mu' d\phi \quad [\text{diffuse (multiple) scattering}]$$


$$+ \frac{\tilde{\omega}}{4\pi} F_0 P(\mu, \phi; -\mu_0, \phi_0) e^{-\frac{\tau}{\mu_0}} \quad [\text{single(direct) scattering}]$$

$$\begin{cases} P: \text{Scattering phase function : redirects } (\mu', \phi') \rightarrow (\mu, \phi) \\ \tilde{\omega} = \frac{\sigma_s}{\sigma_e}: \text{Scattering albedo} \\ \text{scattering cross section/extinction(scattering + absorption) cross section} \end{cases}$$

* remove ϕ dependency using $P(\cos\theta)$ function

* $P, \tilde{\omega}$, Albedo depend on λ , particle size & shape.


$$P(\cos\phi) = \sum_{l=0}^N \tilde{\omega}_l P_l(\cos\phi) \quad : \text{Legendre Polynomial}$$



Radiative transfer equation solver.

→ $\begin{cases} \text{Discrete - ordinates method} \\ \text{Two - Stream and Eddington's approximation} \\ \text{Delta - function adjustment and similarity principle} \\ \delta - \text{Four stream approximation} \end{cases}$

1.3 Terrestrial radiation





$0, \mu$	$\tau_\infty, z_\infty, 0$
τ', μ'	τ', z', P'
τ, μ	τ, z, P
τ', μ'	τ', z', P'
$\tau, 0$	$\tau_s, 0, P_*$

$$F(z) = F^\uparrow(z) - F^\downarrow(z)$$

$$\Delta F = F(z + \Delta z) - F(z)$$

$$\left. \frac{\partial T}{\partial t} \right|_{IR} = - \frac{1}{c_P \rho} \frac{\Delta F}{\Delta P} = - \frac{g}{c_P} \frac{\Delta F}{\Delta u}$$



In spectral bands (monochromatic)


$$\uparrow \mu \frac{dI_\nu(\tau, \mu)}{d\tau} = I_\nu(\tau, \mu) - B_\nu(T) \quad \text{B.C. } \begin{cases} SFC (\tau = \tau_1), I_\nu(\tau, \mu) = B_\nu(T_s) \\ TOP (\tau = 0), I_\nu(0, -\mu) = 0 \end{cases}$$

$$\downarrow -\mu \frac{dI_\nu(\tau, -\mu)}{d\tau} = I_\nu(\tau, -\mu) - B_\nu(T)$$

$$F^\uparrow(\tau) = 2\pi B_\nu(T_s) \int_0^1 e^{-\frac{(\tau_1-\tau)}{\mu}} \mu d\mu + 2 \int_0^1 \int_{\tau_1}^{\tau} \pi B_\nu[T(\tau')] e^{-\frac{(\tau'-\tau)}{\mu}} d\tau' d\mu$$

$$F^\downarrow(\tau) = 2 \int_0^1 \int_0^{\tau} \pi B_\nu[T(\tau')] e^{-\frac{(\tau-\tau')}{\mu}} d\tau' d\mu$$

$$d\tau = -k_\nu \rho dz \quad \tau_1 = \int_0^{u_1} k_\nu du, \quad u_1 = \int_0^\infty \rho dz$$



LW is time consuming !

1.4 Cloud fraction


i) Conventional method (Slingo)

$$f = f_c + f_l$$

f_c : depends on precipitation, p_{top} , p_{bottom}

$$f_l : \text{depends on RH} = 1 - \left[\frac{1 - RH}{1 - RH_0} \right]^{0.5}$$


where RH_0 is the critical value of RH,
which is optimized based on observations



ii) Advanced method (Chou)

- inclusion of ice, liquid
- consistent treatment of water substance for both precipitation & radiative properties.

f_c : uses information of detrainment water substances from sub-grid scale clouds in convective parameterizations



f_l : use the hydrometeor information from microphysics routine, q_c, q_s, q_i, \dots

iii) Radiation properties

$$\tau_i^c = cwp \left[a_i + \frac{b_i}{r_{ei}} \right] f_{ice} \quad (\text{optical thickness})$$

$$w_i^c = 1 - c_i - d_i r_{ei} \quad (\text{co-albedo})$$

$$g_i^c = e_i - f_i r_{ei} \quad (\text{asymmetry factor})$$

$$f_i^c = (g_i^c)^2$$

a-f : coeff : depends upon band and k-

$$\bar{\tau}_c = \sum_i \tau_i \quad i : \text{each gas}$$

(The effective optical thickness for each spectral band)

- The long wave cloud emissivity (E_{cld})

$$c_f' = E_{cld} c_f$$

$$E_{cld} = 1 - e^{-Dk_{abc} cwp}$$

where $D = 1.66$: diffusivity factor

$$k_{abc} = k_l (1 - f_{ice}) + k_i f_{ice} \quad : \text{absorptivity coefficient}$$

- For diagnostic microphysics scheme,

- cloud water scale height


$$h_1 = a \ln(1.0 + \frac{b}{g} \int_{P_t}^{P_s} q dp)$$

- cloud droplet size,

$$r_{ee} \begin{cases} = 10\mu m & \text{over ocean} \\ < 10\mu m & \text{over land} \end{cases} \quad : \text{warm clouds}$$

$$r_{ei} : 10\mu m \text{ (low)} \sim 30\mu m \text{ (high)} \quad : \text{(high) ice clouds}$$

- ice fraction, f_{ice}



- For prognostic microphysics scheme,


: Broadband radiation : Radiative properties are explicitly computed from prognostic water substances

: Simplified radiation : Dudhia (1989)

$$\alpha_p \text{ (absorption coefficient)} = \frac{1.66}{2000} \left(\frac{\pi N_0}{\rho_{rs}^3} \right)^{\frac{1}{4}} \quad m^2 g^{-1} = \begin{cases} 2.34 \times 10^{-3} \quad m^2 g^{-1} & \text{for snow} \\ 0.33 \times 10^{-3} \quad m^2 g^{-1} & \text{for rain} \end{cases}$$

$$u_p \text{ (effective water path length)} = \left(\rho q_{rs} \right)^{\frac{3}{4}} \Delta z \times 1000 \quad gm^{-2} \rightarrow \tau_p \text{ (transmission)} = \exp(-\alpha_p u_p)$$

iv) Cloud overlapping



Maximum overlapping : 0.4

Minimum overlapping : 1.0

$$\text{Random overlapping} : H + (1-H)M + \{1-H-(1-H)M\}L = 0.6$$

- Computation :

τ is scaled by A_c (cloud cover) at a given layer.





- Flux for each of $A_c, (1-A_c)$ $\xrightarrow{A_{free}, A_{cld}}$ summation

- Issues : $A_c = 0$ or 1 \leftarrow in WRF versus partial cloudiness in GFS



2. Land-surface processes

2.1 Concept : Surface layer + soil model



Atmospheric surface layer : the lowest part of the atmospheric boundary layer (typically about a tenth of the height of the BL) where mechanical (shear) generation of turbulence exceeds buoyant generation or consumption. Turbulent fluxes and stress are nearly constant with height.

In atmospheric models, it is defined the height of the lowest model level.

2.2 Surface layer parameterization

Surface layer schemes calculate friction velocities and exchange coefficients that enable the calculation of Surface heat and moisture fluxes by the land-surface models. These fluxes provide a lower boundary condition for the vertical transport done in the PBL Schemes.

Over water surfaces, the surface fluxes and surface diagnostic fields are computed in the surface layer scheme itself. Sea surface temperature can be predicted by the surface energy budget and mixed layer mixing

1) Bulk method

$$H_0 = \rho C_p C_H |\vec{V}_a| \Delta T$$

$$E_0 = \rho L C_H |\vec{V}_a| \Delta q M_a$$


$$\vec{\tau}_0 = \rho C_D |\vec{V}_a| \vec{V}_a$$

C_D, C_H : prescribed
 $= 0.01$ over land, $= 0.001$ over water

2) Monin-Obukov similarity

$$\frac{k_z}{u_*} \frac{\partial u}{\partial z} = \phi_m(z/L), \quad \frac{k_z}{u_*} \frac{\partial \theta}{\partial z} = \phi_t(z/L)$$

$$\text{Integrate, } F_m = \int_{z_0}^{h_s} \frac{dz}{z} \phi_m dz = \ln\left(\frac{h_s}{z_0}\right) - \psi_m(h_s, z_0, L)$$



* **Profile function** : ϕ_m and ϕ_t

Dyer and Hicks formula for similarity

- unstable ($L < 0$)



$$\phi_m = \left(1 - 16 \frac{0.1h}{L}\right)^{-\frac{1}{4}} \quad \text{for } u, v$$

$$\phi_t = \left(1 - 16 \frac{0.1h}{L}\right)^{-\frac{1}{2}} \quad \text{for } \theta, q$$

- stable ($L > 0$)

$$\phi_m = \phi_t = \left(1 + 5 \frac{0.1h}{L}\right)$$

$$\text{where } L = u_*^2 \bar{\theta} / (kg\theta_*) = -\frac{\rho C_p \theta_0 u_*^3}{kgH_0}$$



$$\frac{h_s}{L} = \frac{\phi_m^2 (hs/L)}{\phi_t^2 (hs/L)} Ri = \zeta \text{ (Zeta)}$$

$$\text{where } Ri = \frac{g}{\theta} \frac{\partial \theta}{\partial z} \quad \left(\frac{\partial u}{\partial z}\right)^{-2}$$

* **Useful relation** :

Given the $F_m, F_H, C_D = k^2 / F_m^2, C_Q = C_H = k^2 / (F_m F_t), u_* = kU / F_m$

$$\tau_0 = \rho k_m \frac{du}{dz} = -\overline{u'w'} = \rho C_p U^2$$

$$H_0 = -\rho C_p k_h \frac{d\theta}{dz} = \rho C_p \overline{\theta'w'} = -\rho C_p C_H U \Delta \theta$$

$$E_0 = -\rho L \overline{q'w'} = -\rho L C_q U \Delta q$$

2.3 Soil model

1) Slab model : force-restore method

$$\frac{\partial T_s}{\partial t} = \lambda_T (R_n - LE - H) - \frac{2\pi}{\tau} (T_s - T_a)$$

$$\frac{\partial T_m}{\partial t} = \frac{1}{\tau} (T_s - T_m) : T_m, \text{ daily mean}$$



2) Multi-layer model : OSU method

$$\frac{\partial T_s}{\partial t} = \lambda_T (R_n - LE - H) : \text{surface T}$$

$$(\rho C)_i \frac{\partial T_g}{\partial t} = \frac{\partial}{\partial z} (\lambda T_g \frac{\partial T_g}{\partial z}) : \text{soil T}$$

$$\frac{\partial \Theta}{\partial t} = \frac{\partial}{\partial z} (D \frac{\partial \Theta}{\partial z}) + K \frac{\partial \Theta}{\partial z} + F_\Theta : \text{soil moisture}$$


- NOAH, SIB, PLACE, VIC, CLM, etc



2.4 Vegetation type → z_0 , Albedo

Vegetation types


MODIS (satellite)




13 type data set
1 degree

12 type data set
20 min

The Simple Biosphere model (SiB)




United States Geological Survey's (USGS)



Albedo and Roughness length (z_0)


Albedo

Sib




16.6 %↓

USGS




25.5 %

USGS-Sib




-20 -10 -5 5 10 20


Roughness
length



60.8cm↓



38.0cm↓




-80 -40 -20 20 40 80

3. Vertical diffusion (PBL)


3.1 Concept

- computes the parameterized effects of vertical turbulent eddy diffusion of momentum, water vapor and sensible heat fluxes




3.2 Planetary Boundary Layer Structure : schematic

Daytime profiles



Daytime flux profiles



3.3 Classifications : how to determine, k_c

i) Local diffusion (Louis 1979)

$$\frac{\partial c}{\partial t} = \frac{\partial}{\partial z} (-\bar{w}\bar{c}) = \frac{\partial}{\partial z} (k_c \frac{\partial c}{\partial z}) \quad k_c : \text{diffusivity}, \quad k_m, k_t = l^2 f_{m,t}(Ri) \left| \frac{\partial U}{\partial z} \right|$$


Local Richardson number

ii) Nonlocal diffusion (Troen and Mahrt 1986)

$$\frac{\partial c}{\partial t} = \frac{\partial}{\partial z} (-\bar{w}\bar{c}) = \frac{\partial}{\partial z} (k_c (\frac{\partial c}{\partial z} - \gamma_c))$$

$$k_{zm} = k_w z (1 - \frac{z}{h})^p, \quad h = R_{ibcr} \frac{\theta_m}{g} \frac{U^2(h)}{(\theta_v(h) - \theta_s)}$$

$$\theta_s = \theta_{va} + \theta_T (= b \frac{(\theta_v' w')_0}{w_s}), \quad w_s = u_* \phi_m^{-1}$$



iii) Eddy mass-flux diffusion (Siebesma and Teixiera 2000)

$$\frac{\partial c}{\partial t} = \frac{\partial}{\partial z} (-\bar{w}\bar{c}) = -\frac{\partial}{\partial z} \left[-k_c \frac{\partial \bar{c}}{\partial z} + M(c_u - \bar{c}) \right]$$

small eddies strong updrafts


iv) TKE (Turbulent Kinetic Energy) diffusion (Mellor and Yamada 1982)

TKE equation :

$$\frac{\overline{\partial u_i u_j}}{\partial t} + \overline{u_j} \frac{\partial \overline{u_i u_j}}{\partial x_j} = - \frac{\partial}{\partial x_k} \left[\overline{u_i u_j u_k} + \frac{1}{\rho} \dots \right]$$

$$\overline{u_i u_j} \implies k_z = \text{fn (TKE)}$$

3.4 Local versus nonlocal



Local scheme typically produces unstable mixed layer


Hong and Pan (1996)

4. Gravity Wave Drag


- GWDO : GWD induced by sub-grid scale orography
- GWDC : GWD induced by precipitating deep convection

4.1 Concept

This scheme (GWDO) includes the effect of mountain induced gravity wave drag from sub-grid scale orography including convective breaking, shear breaking and the presence of critical levels. Effects are strong in the presence of strong vertical wind shear and thermally stable layer.



4.2 Enhanced lower tropospheric gravity wave drag (Kim and Arakawa 1995)



Stress at reference level :

$$\tau_0 = -E \frac{1}{\Delta x} \frac{\rho_0 U_0^3}{N_0} \frac{Fr^2}{Fr^2 + 0.5/OC}, \quad U_0 = \frac{1}{h} \int_{k=1}^{k=k_{pbl}} U dz$$

Reference level (KA95) : Max (2, KPBL)


Conventional : the conventional Ri number-based wave- breaking mechanism using the saturation hypothesis, which works mainly in the upper atmosphere

Advanced: the new orographic statistics-based wave- breaking mechanism using half-theory (Scorer parameter $\sim BVF^{**2} / U^{**2}$) and half-empiricism obtained from mesoscale mountain wave simulations, which works mainly in the lower atmosphere. Flow blocking is also introduced recently.

4.3 Impact of GWDO


Zonal-averaged zonal wind (96/97 DJF)

RA2




Contour : Zonal averaged
zonal wind
Shaded: Deviations from
the RA2

NOGWD



GWD




Kim and Arakawa
→Improves upper level jets
→Improves the sea level
pressure

(Kim and Hong, 2009)

4.4 Convective GWD (CWDC)


Columnar CGWD parameterizations

- **Chun and Baik (1998, 2002):** The momentum flux spectrum for the CGWD parameterization was first analytically formulated
- **Chun et al. (2008):** A nonlinear source effect was included in the CGWD parameterization of Song and Chun (2005) that had taken account of a diabatic source alone



Ray-based CGWD parameterization


- **Song and Chun (2008):** GW propagation properties were explicitly calculated and a three-dimensional propagation of GWs was realistically represented
- **Choi and Chun (2011):** Two free parameters, the moving speed of the convective source and wave-propagation direction, were determined




Prof. Chun, Hye-Young, chunhy@yonsei.ac.kr

4.5 Improvement by GWDC (Jeon et al. 2010, APJAS)


a) SAS_CB98 Zonal wind difference




a) SAS_CB98 Temperature difference



c)



b)



Precipitation Processes

Concept : precipitation algorithms (CPS and MPS)


- In real atmosphere, dynamical motion $\rightarrow RH > 1 \Rightarrow$ clouds form \rightarrow produces rain
- In modeled atmosphere, $RH < 1$

But generate clouds by releasing CAPE \rightarrow requires parameterized process

Deep convection : 1~10km



$$\overline{RH} < 1$$



$\Delta x \implies 0$, more grid-resolvable precipitation



Thus, we need the cumulus parameterization scheme to account for releasing conditional instability due to subgrid scale motion

- Grid-resolvable (MPS): Supersaturation \rightarrow clouds
- Subgrid scale (CPS) : CAPE removal \rightarrow clouds


5. Deep Convection

Parameterized convection
Cumulus convection
Subgridscale precipitation
Implicit precipitation

5.1 Concept

- represents deep precipitating convection and feedback to large-scale
- must formulate the collective effects of subgridscale clouds in terms of the prognostic variable of grid scale

● Closure assumption



Jul 25, 2010, Seoul




b

Processes CP Schemes Need to Account For

Convective Initiation

Deep Convection




Shallow Convection at Top of Mixed Layer



Vertical Heating / Cooling


Vertical Drying / Moistening

Precipitation




The COMET Program

CPS does not consider the detailed evolution of convection !



Charleston SC 22 Apr 1998 2200 UTC

(dashed=model, darker red/green=1st guess BMJ,
lighter red/green=enthalpy adjusted BMJ profiles)



post-convective adjustment, 1h later

Michael Baldwin

CPS consider changes in profile before and after convection

5.2 Kuo scheme (1965)

: Cloud is formed in proportional to column-integrated moisture convergence

$$M_t = -\frac{1}{g} \int_0^{P_s} \nabla \cdot (vq) dp + F_{g_s}$$

$$\int_0^{P_s} \frac{\partial q}{\partial t} dp = gbM_t, \quad b : \text{moistening factor}$$

$$\int_0^{P_s} \theta_c dp = gL(1-b)M_t \quad \frac{\theta_c}{\pi} = \frac{\theta_a - \theta}{\tau}$$

- Modified Kuo scheme

Krishnamurti et al. (1980, 1983) : proportional to

vertical advection of moisture

$$M_t = -\frac{1}{g} \left\{ \int_0^{P_s} \omega \frac{\partial q}{\partial p} dp + F_{q_s} \right\}$$

Anthes (AK : 1977) : a revised moistening factor
and parcel buoyancy


$$b = \left(\frac{1 - RH}{1 - RH_c} \right)^n$$

- Heating and moistening profiles

$$\frac{d\theta}{dt} = \frac{1}{\pi} [gL(1-b)M_t Q_1 + Q_r]$$

$$\frac{dq}{dt} = -g(1-b)M_t Q_2$$


$$\int_0^{P_s} Q_1 dp = \int_0^{P_s} Q_2 dp = 1$$



Remarks : Kuo scheme produces excessive point maximum precipitation. Also, warm bias in the lower troposphere. This scheme take into account of large-scale mixing as well as turbulent mixing

5.3 Betts-Miller scheme (1986)

: adjust toward reference profiles that are based on observational evidence of convective equilibrium



Reference profile to be adjusted : originally based on tropical cyclones

$$\theta'_R(P) = \bar{\theta}(P_B) + \beta M_\theta (P - P_B)$$

$$\frac{\partial q}{\partial p} = \beta \left(\frac{\partial q}{\partial p^*} \right)_M \quad \beta = \frac{\partial p^*}{\partial p} \quad p^* : \text{saturation pressure} \quad (= 1.2 \text{ for example})$$

$$M_\theta = 0.85 \left(\frac{\partial \theta^*}{\partial P^*} \right)_M \quad M : \text{Mixing line}$$

$$\text{Energy Constraints : } \int_{P_B}^{P_{T+1}} C_P (T_R - \bar{T}) dp = \int_{P_B}^{P_{T+1}} (q_R - \bar{q}) dp = 0$$

- Convective tendencies & Precipitation


$$\left(\frac{\partial \bar{T}}{\partial t} \right)_{Cu} = \frac{T_R - \bar{T}}{\tau}$$

$$\left(\frac{\partial \bar{q}}{\partial t} \right)_{Cu} = \frac{q_R - \bar{q}}{\tau}$$

$$Precip = \int_{P_0}^{P_T} \left(\frac{q_R - \bar{q}}{\tau} \right) \frac{dP}{g} = - \frac{C_p}{L} \int_{P_0}^{P_T} \left(\frac{T_R - \bar{T}}{\tau} \right) \frac{dp}{g}$$

- Adjustment scheme :

- Manabe (hard adjustment : toward a moist adiabat)
- Kuo, BM (soft adjustment : toward reference profiles)



- Remarks :

- Uncertainty in reference profiles → 2×CO₂ → similar climate ???


5.4 Mass-flux schemes : Arakawa-Schubert (1974)

i) Concept

-- Mass flux approach, cloud ensemble, quasi-equilibrium

-- Theoretical frame work for CPS

- Area is large enough so that cloud ensemble can be a statistical entity
- Area is small enough so that cloud environment is approximately uniform horizontally



M_i : vertical mass flux through ith cloud

σ_i : fractional area covered by ith cloud

$M_c \equiv \sum_i M_i$: total vertical mass flux

$$\rho M = M_c + \tilde{M}$$

environment

: net mass flux/unit large-scale horizontal area

ii) Quasi-equilibrium : cloud forcing ~ large-scale adjustment

- CPS computes the warming (cooling) in the grid box due to adiabatic descent (ascent), rather than computing latent heat releaser in cloud models




FIG. 4.10. Model for updraft and downdraft of cloud type λ (from Johnson 1976).

iii) Energy budget equations

Large-scale flux across grid box Exchange of S between environment and clouds

$$\frac{\partial}{\partial t} \rho (1 - \sigma_c) \tilde{S} = -\vec{\nabla} \cdot (\overline{\rho \tilde{V} S}) - \frac{\partial}{\partial Z} (\tilde{M} \tilde{S}) - \sum_i \left(\frac{\partial M_i}{\partial Z} + \rho \frac{\partial \sigma_i}{\partial t} \right) S_{ib} - LE + \tilde{Q}_R$$

$$\frac{\partial}{\partial t} \rho \sum_i \sigma_i S_i = -\frac{\partial}{\partial z} \left(\sum_i M_i S_i \right) + \sum_i \left(\frac{\partial M_i}{\partial z} + \rho \frac{\partial \sigma_i}{\partial t} \right) S_{ib} + \sum_x (LC_x + Q_{Ri})$$

$S_i : C_p T + gz$ (dry static energy) of i^{th} cloud

$S_{ib} : C_p T + gz$ of the air entraining into or detrainning from the i^{th} cloud

C_i : condensation in the i^{th} cloud

E : evaporation of liquid water in the environment

Q_r : Radiative heating

- Entrainment : $\frac{\partial M_i}{\partial z} + \rho \frac{\partial \sigma_i}{\partial t} > 0, \quad S_{ib} = \tilde{S}$

- Detrainment : $\frac{\partial M_i}{\partial z} + \rho \frac{\partial \sigma_i}{\partial t} < 0, \quad S_{ib} = S_i$

iv) Approximation

- Assume $\sigma_c \ll 1$, $\bar{s} \approx \tilde{s}$

$$\begin{aligned}\frac{\partial}{\partial t} \rho \bar{s} &= -\nabla \cdot (\rho \bar{v} \bar{s}) - \frac{\partial}{\partial z} (\rho \bar{w} \bar{s}) - \overline{\nabla \cdot (\rho \bar{v} s - \rho \bar{v} \bar{s})} \\ &\quad + M_c \frac{\partial \bar{s}}{\partial z} - \sum_{dc} \left(\frac{\partial M_i}{\partial z} + \rho \frac{\partial \sigma_i}{\partial t} \right) (\delta_i - \bar{s}) - LE + \dot{\theta}_R\end{aligned}$$

Detraining clouds dc

detrainment, entrainment

Adiabatic warming due to hypothetical
subsidence between the clouds

$$\begin{aligned}\frac{\partial}{\partial t} \rho \bar{q} &= -\nabla \cdot (\rho \bar{v} \bar{q}) - \frac{\partial}{\partial z} (\rho \bar{w} \bar{q}) - \overline{\nabla \cdot (\rho \bar{v} q - \rho \bar{v} \bar{q})} \\ &\quad + M_c \frac{\partial \bar{q}}{\partial z} - \sum_{dc} \left(\frac{\partial M_i}{\partial z} + \rho \frac{\partial \sigma_i}{\partial t} \right) (q_i - \bar{q}) - E\end{aligned}$$

- Spectral cloud ensemble :

$$M_c(z) = \int_0^{\lambda_{\max}} m(z, \lambda) d\lambda \quad \text{Sub-ensemble}$$

mass flux of between λ and $d\lambda + \lambda$

$$= \int_0^{\lambda_{\max}} m_B(\lambda) \eta(z, \lambda) d\lambda$$

Mass flux at cloud base

$$\eta(z, \lambda) \equiv \frac{m(z, \lambda)}{m_B(\lambda)} \quad ; \quad \text{normalized subensemble mass flux}$$

v) Closure

$$\frac{\partial m(z, \lambda)}{\partial z} = \mu(z, \lambda) \eta(z, \lambda)$$

$$\eta(z, \lambda) = e^{\lambda(z - z_B)} ; \text{ mass flux profile}$$

- Cloud work function

$$A(\lambda) = \int_{z_B}^{z_D(\lambda)} \eta(z, \lambda) g \frac{T_c(z, \lambda) - \bar{T}(z)}{\bar{T}} dz$$

- Q-G equilibrium

$$\frac{dA(\lambda)}{dt} = \left. \frac{dA(\lambda)}{dt} \right|_{LS} + \left. \frac{dA(\lambda)}{dt} \right|_C ; 0$$


Large-scale forcing
 >0 : destabilized

Adjustment
 <0 : stabilization

Kernel : Cloud scheme kinetic energy

$$K_{ij} = \frac{A'_i - A_i}{(m_B \Delta t)} \quad \sum_j^{i_{\max}} K_{ij} (m_B \Delta t)_j + F_i = 0 \Rightarrow m_B$$

————— compute $\frac{\partial \bar{s}}{\partial t}, \frac{\partial \bar{q}}{\partial t}$ with η, m_B



5.5 Other schemes

* AS type mass flux scheme

Grell scheme (1993) : Updraft/downdraft couplet without lateral mixing to find the deepest cloud

Simplified AS (SAS, Han and Pan 2011): revised cloud physics from the Grell

Relaxed AS (RAS, Moorthi and Suarez 1992): linearized profile function

* Other mass flux schemes : Low-level control convective schemes (Stensrud 2007)

Kain and Fritsch (1993) : CAPE based sophisticated convective plume model

Emanuel (1991) : Stochastic mixing cloud model

Tiedtke (1989) : Large-scale moisture convergence (KUO) based mass flux

Gregory-Rowntree (1990): Parcel buoyancy based turbulence in cloud model

6. Shallow Convection

6.1 Concept

more vigorous vertical mixing of q and T above the mixed layer top. With the enhanced vertical eddy transport between LCL and inversion level, this process does not allow the excess moisture trapped near the surface in synoptically inactive regions (**non-precipitating convection**).

- Cooling and moistening above LCL and heating and warming below.



6.2 Classification

- Moist adjustment type : Betts and Miller (1993), Lock et al. (2000), Tiedtke (1983)
- Mass flux type : Kain (2004), Park and Bretherton (2009), Han and Pan (2011)

Tiedtke (1983)

$$\frac{\partial T}{\partial t} = \frac{1}{\rho} \frac{\partial}{\partial z} \left(\rho K \left[\frac{\partial T}{\partial z} + \Gamma \right] \right)$$

$$\frac{\partial q}{\partial t} = \frac{1}{\rho} \frac{\partial}{\partial z} \left(\rho K \frac{\partial q}{\partial z} \right)$$

Han and Pan (2011)

$$\frac{1}{\eta} \frac{\partial \eta}{\partial z} = \varepsilon - \delta$$

$$\frac{\partial(\eta s)}{\partial z} = (\varepsilon \bar{s} - \delta s) \eta$$


$$\frac{\partial[\eta(q_v + q_l)]}{\partial z} = \eta[\varepsilon \bar{q}_v - \delta(q_v + q_l) - r]$$

6.3 Impact of the shallow convection scheme


JJA 1996 simulation in a GCM

Temperature


NO - RA2






RH



Tiedtke - NO



Han and Pan - NO



dry
moist

mixing

mixing

7. Non-convective Precipitation

large-scale precipitation
grid-resolvable scale precipitation
explicit moisture scheme
cloud scheme
microphysics scheme

7.1 Concept

- Remove supersaturation after deep and shallow convection, and feedback to large-scale

7.2 Classification : according to the complexity in microphysics

i) Diagnostic : condensation, evaporation of falling precipitation

ii) Bulk microphysics :

hydrometeors with size distribution in inverse-exponential function

- Single moment : predict mixing ratios of hydrometeors
- Double moment : + number concentrations
- Triple moment : + reflectivity

iii) Bin microphysics : divides the particle distribution into a number of finite size or mass categories.

7.3 Precipitate size distributions

Marshall and Palmer(1948) : exponential law

Heymsfield and Platt (1984) : Power law

$$N_R(D_R) = aD_R^{-b}$$

The rain and snow particles are assumed to follow the size distribution derived by Marshall and Palmer(1948), and Gunn and Marshall(1958), respectively. The size distributions for both rain and snow are formulated according to an inverse-exponential distribution and its formula for rain can be expressed by

$$N_R(D_R) = N_{0R} \exp(-\lambda_R D_R)$$

for rain, where N_{0R} is the intercept parameter of the rain distributions.

The slope parameter of the size distributions for rain λ_R is determined by multiplying (A1) by drop mass (A4) and integrating over all diameters and equating the resulting quantities to the appropriate water contents ($= \rho q_R$). This may be written as,

$$\lambda_R = \left(\frac{\pi \rho_w N_{0R}}{\rho q_R} \right)^{1/4}$$



$$\Gamma(x) = \int_0^{\infty} t^{x-1} \exp(-t) dt$$

$$\int_0^{\infty} D_R^{4-1} \exp(-\lambda_R D_R) dD_R \\ = \Gamma(4) / \lambda_R^4$$

7.4 Bulk Method : 1-Moment versus 2-Moment

Mixing Ratio

(1-moment/ 2-moment scheme)

$$\left(\int \frac{dM(D_R)}{dt} dN_{DR} \right) / \rho = \frac{dq}{dt} (kg kg^{-1} s^{-1})$$

Number concentration

(2-moment scheme)

$$\left(\int \frac{d \text{Prob}(D_R)}{dt} dN_{DR} \right) = \frac{dN}{dt} (m^{-3} s^{-1})$$




FIG. 9.21. Evolution of a drop spectrum in a subsaturated rainshaft including the effects of coalescence and breakup. The spectrum at the top of the rainshaft is Marshall-Palmer with a rainfall rate of 110 mm h^{-1} . (.....), 0 km; (---), 1 km; (—), 2 km; $R = 110 \text{ mm h}^{-1}$; $t = 24 \text{ min}$. From Tzivivon et al. (1989). *Journal of Atmospheric Sciences*, 46, 21. American Meteorological Society. Reproduced with permission.

Single moment scheme

$$dN_{DR} = N_{0R} \exp(-\lambda_R D_R) dD_R$$


Double moment scheme

$$dN_{DR} = N_R \lambda_R^2 (N_R) D_R \exp(-\lambda_R D_R) dD_R$$

7.5 Bulk Method : 1-Moment (WSM) versus 2-Moment (WDM)

Cold rain processes :


(Hong et al. 2004; Hong and Lim 2006)



q for 4 hydrometeors will be predicted (Single Moment)

Warm rain processes :

(Khairoutdinov and Kogan 2000; Cahard and Pinty 2000)



N, q for 2 hydrometeors will be predicted (Double Moment)

N: Cloud water, Rain, CCN

Q: Cloud water, Rain, Ice, Snow, Graupel, Vapor

WDM6

(Lim and Hong, 2010)

7.6 WDM6 code

** Warm rain processes (Hong and Lim 2010)

```

| warm rain processes
|
| - follows the double-moment processes in Lim and Hong
| =====
|
| do k = kts, kte
|   do i = its, ite
|     supsat = max(q(i,k),qmin)-qs(i,k,1)
|     satdt = supsat/dtcld
|
|   praut: auto conversion rate from cloud to rain [LH 9] [CP 17]
|   (QC->QR)
|
|     lencon = 2.7e-2*den(i,k)*qci(i,k,1)*(1.e20/16.*rslope2(i,k)
|           *rslope2(i,k)-0.4)
|     lenconcr = max(1.2*lencon, qcrmin)
|     if(avedia(i,k,1).gt.di15) then
|       taucon = 3.7/den(i,k)/qci(i,k,1)/(0.5e6*rslope2(i,k)-7.5)
|       praut(i,k) = lencon/taucon
|       praut(i,k) = min(max(praut(i,k),0.),qci(i,k,1)/dtcld)
|
|   nraut: auto conversion rate from cloud to rain [LH A6] [CP 18 & 19]
|   (NC->NR)
|
|     nraut(i,k) = 3.5e9*den(i,k)*praut(i,k)
|     if(qrs(i,k,1).gt.lenconcr)
|       nraut(i,k) = ncr(i,k,3)/qrs(i,k,1)*praut(i,k)
|       nraut(i,k) = min(nraut(i,k),ncr(i,k,2)/dtcld)
|     endif
|
|   pracw: accretion of cloud water by rain      [LH 10] [CP 22 & 23]
|   (QC->QR)
|   nracw: accretion of cloud water by rain      [LH A9]
|   (NC->)
|
|     if(qrs(i,k,1).ge.lenconcr) then
|       if(avedia(i,k,2).ge.di100) then
|         nracw(i,k) = min(ncrk1*ncr(i,k,2)*ncr(i,k,3)*(rslope3(i,k)
|           + 24.*rslope3(i,k,1)),ncr(i,k,2)/dtcld)
|         pracw(i,k) = min(pi/6.*(denr/den(i,k))*ncrk1*ncr(i,k,2)
|           *ncr(i,k,3)*rslope3(i,k)*(2.*rslope3(i,k)
|           + 24.*rslope3(i,k,1)),qci(i,k,1)/dtcld)
|       else
|         nracw(i,k) = min(ncrk2*ncr(i,k,2)*ncr(i,k,3)*(2.*rslope3(i,k)
|           *rslope3(i,k,1)+5040.*rslope3(i,k,1)
|           *rslope3(i,k,1)),ncr(i,k,2)/dtcld)
|         pracw(i,k) = min(pi/6.*(denr/den(i,k))*ncrk2*ncr(i,k,2)
|           *ncr(i,k,3)*rslope3(i,k)*(6.*rslope3(i,k)
|           *rslope3(i,k,1)+5040.*rslope3(i,k,1)*rslope3(i,k,1)
|           ,qci(i,k,1)/dtcld)
|       endif
|     endif
|   endif
| 
```

*Auto conversion from cloud to rain [C → R]

$$\text{Praut} [\text{kg kg}^{-1} \text{s}^{-1}] = L / \tau \quad L = 2.7 \times 10^{-2} \rho_a q_c \left(\frac{10^{20}}{16 \lambda_c^4} - 0.4 \right)$$

$$\tau = 3.7 \frac{1}{\rho_a q_c} \left(\frac{0.5 \times 10^6}{\lambda_c} - 7.5 \right)^{-1}$$

$$\text{Nraut} [\text{m}^{-3} \text{s}^{-1}] = 3.5 \times 10^9 \frac{\rho_a L}{\tau}$$

*Accretion of cloud water by rain [C → R]

$$D_R \geq 100 \mu\text{m}$$

$$\text{Pracw} [\text{kg kg}^{-1} \text{s}^{-1}] = \frac{\pi}{6} \frac{\rho_w}{\rho_a} K_1 \frac{N_C N_R}{\lambda_c^3} \left\{ \frac{2}{\lambda_c^3} + \frac{24}{\lambda_R^3} \right\}$$

$$\text{Nracw} [\text{m}^{-3} \text{s}^{-1}] = -K_1 N_C N_R \left\{ \frac{1}{\lambda_c^3} + \frac{24}{\lambda_R^3} \right\}$$

$$D_R < 100 \mu\text{m}$$

$$\text{Pracw} [\text{kg kg}^{-1} \text{s}^{-1}] = \frac{\pi}{6} \frac{\rho_w}{\rho_a} K_2 \frac{N_C N_R}{\lambda_c^3} \left\{ \frac{6}{\lambda_c^6} + \frac{5040}{\lambda_R^6} \right\}$$

$$\text{Nracw} [\text{m}^{-3} \text{s}^{-1}] = -K_2 N_C N_R \left\{ \frac{2}{\lambda_c^6} + \frac{5040}{\lambda_R^6} \right\}$$


Resolution Dependency

Cut-off horizontal grid length for parameterizations


- PBL : ~50 m (Mirocha, 2008 WRF workshop)
- GWDO : ~ 3 km (hydrostatic approximation)
- GWDC: ~ 3 km (go with CP)
- Cumulus parameterization : ~ 3 km (cloud resolving scale)

Gray-zone : partly resolved and partly parameterized


- CPS (1 km~10 km) : Gerard, Grell and Freitas, Arakawa and Ming, Pan and Han,
- PBL (100 m~1 km) : Honnert, Boulle, Shin and Hong



OBS



No CPS



With CPS

References

- Anthes, R. A., 1977: A cumulus parameterization scheme utilizing a one-dimensional cloud model. *Mon. Wea. Rev.*, **105**, 270–286.
- Arakawa, A., and W. H. Schubert, 1974: Interaction of a cumulus cloud ensemble with the large-scale environment, Part I. *J. Atmos. Sci.*, **31**, 674-701.
- Betts, A. K., and M. J. Miller, 1986: A new convective adjustment scheme. Part II: Single column tests using GATE wave, BOMEX, ATEX and arctic air-mass data sets. *Quart. J. Roy. Meteor. Soc.*, **112**, 693-709.
- Betts, A. K., and M. J. Miller, 1993: The Betts-Miller scheme. *The representation of cumulus convection in numerical models*, K. A. Emanuel and D. J. Raymond, Eds., Amer. Meteor. Soc., 246 pp.
- Chen, F., and J. Dudhia, 2001: Coupling an advanced land-surface/hydrology model with the Penn State/NCAR MM5 modeling system. Part I: Model description and implementation. *Mon. Wea. Rev.*, **129**, 569-585.
- Chun, H.-Y., and J.-J. Baik, 1998: Momentum flux by thermally induced internal gravity waves and its approximation for large-scale models. *Journal of the Atmospheric Sciences*, **55**, 3299-3310.
- Cohard, J.-M., and J.-P. Pinty, 2000: A comprehensive two-moment warm microphysical bulk scheme. I: Description and tests. *Quart. J. Roy. Meteor. Soc.*, **126**, 1815–1842.
- Dyer, A. J., and B. B. Hicks, 1970: Flux-gradient relationships in the constant flux layer. *Quart. J. Roy. Meteor. Soc.*, **96**, 715-721.
- Emanuel, K. A., 1991: A scheme for representing cumulus convection in large-scale models. *J. Atmos. Sci.*, **48**, 2123–2335.
- Gregory, D., and P. R. Rowntree, 1990: A mass flux convection scheme with representation of cloud ensemble characteristics and stability-dependent closure. *Mon. Wea. Rev.*, **118**, 1483-1506.
- Grell, G. A., 1993: Prognostic evaluation of assumptions used by cumulus parameterizations. *Mon. Wea. Rev.*, **121**, 764-787.

References

- Han, J., and H.-L. Pan, 2011: Revision of convection and vertical diffusion schemes in the NCEP global forecast system. *Wea. and Forecasting*, **26**, 520–533.
- Heymsfield, A. J., and C. M. R. Platt, 1984: A parameterization of the particle size spectrum of ice clouds in terms of the ambient temperature and the ice water content. *J. Atmos. Sci.*, **41**, 846–855.
- Hong, S.-Y., and H.-L. Pan, 1996: Nonlocal boundary layer vertical diffusion in a Medium-Range Forecast model. *Mon. Wea. Rev.*, **124**, 2322–2339.
- Hong, S.-Y., J. Dudhia, and S.-H. Chen, 2004: A revised approach to ice microphysical processes for the bulk parameterization of clouds and precipitation. *Mon. Wea. Rev.*, **132**, 103–120.
- Hong, S.-Y., and J.-O. J. Lim, 2006: The WRF single-moment 6-class microphysics scheme (WSM6). *J. Korean Meteor. Soc.*, **42**, 129–151.
- Kain, J. S., and J. M. Fritsch, 1993: Convective parameterization for mesoscale models: The Kain-Fritsch scheme, *The representation of cumulus convection in numerical models*, K. A. Emanuel and D. J. Raymond, Eds., Amer. Meteor. Soc., 246 pp.
- Kain, J. S., 2004: The Kain-Fritsch convective parameterization : An update. *J. Appl. Meteor*, **43**, 170–181.
- Khairoutdinov, M., and Y. Kogan, 2000: A new cloud physics parameterization in a large-eddy simulation model of marine stratocumulus. *Mon. Wea. Rev.*, **128**, 229–243.
- Kim, Y.-J., and A. Arakawa, 1995: Improvement of orographic gravity-wave parameterization using a mesoscale gravity-wave model. *J. Atmos. Sci.*, **52**, 1875–1902.
- Kim, Y.-J., and S.-Y. Hong, 2009: Interaction between the orography-induced gravity wave drag and boundary layer processes in a global atmospheric model. *Geophys. Res. Lett.*, **36**, L12809, doi: 10.1029/2008GL037146.

References

- Kuo, H.-L., 1965: On the formation and intensification of tropical cyclones through latent heat release by cumulus convection. *J. Atmos. Sci.*, **22**, 40–63.
- Krishnamurti, T. N., Y. Ramanathan, H. L. Pan, R. J. Pasch, and J. Molinari, 1980: Cumulus parameterization and rainfall rates I. *Mon. Wea. Rev.*, **108**, 465–472.
- Krishnamurti, T. N., L.-N. Simon, and R. J. Pasch, 1983: Cumulus parameterization and rainfall rates I. *Mon. Wea. Rev.*, **111**, 815–4828.
- Lim, K.-S. S., and S.-Y. Hong, 2010: Development of an effective double-moment cloud microphysics scheme with prognostic cloud condensation nuclei (CCN) for weather and climate models. *Mon. Wea. Rev.*, **138**, 1587–1612.
- Lock, A. P., A. R. Brown, M. R. Bush, G. M. Martin, and R. N. B. Smith, 2000: A new boundary layer mixing scheme. Part I: Scheme description and single-column model tests. *Mon. Wea. Rev.*, **128**, 3187–3199.
- Marshall, J. S., and W. MaK. Palmer, 1948: The distribution of raindrops with size. *J. Meteor*, **5**, 165–166.
- Mellor, G., and T. Yamada, 1982: Development of a turbulence closure model for geophysical fluid problems. *Rev. Geophys. Space Phys.*, **20**, 851–875.
- Mirocha, J. D., J. K. Lundquist, F. K. Chow, and K. A. Lundquist, 2008: Demonstration of an improved subgrid stress closure for WRF. *The 8th WRF user's workshop*, Boulder, CO, NCAR, 5-4.
- Monin, A. S., and A. M. Obukhov, 1954: Basic laws of turbulent mixing in the surface layer of the atmosphere. *Trudy Geofiz. Inst. Akad. Nauk SSSR*, **24**, 163–187.

References

- Moorthi, S., and M. J. Suarez, 1992: Relaxed Arakawa-Schubert : A parameterization of moist convection for general circulation models. *Mon. Wea. Rev.*, **120**, 978-1002.
- Park, S., and C. S. Bretherton, 2009: The University of Washington shallow convection and moist turbulence schemes and their impact on climate simulations with the Community Atmosphere Model. *J. Climate*, **22**, 3449–3469.
- Shin, H., and S.-Y. Hong, 2009: Quantitative precipitation forecast experiments of heavy rainfall over Jeju Island on 14-16 September 2007 using the WRF model. *Asia-Pacific J. Atmos. Sci.*, **45**, 71-89.
- Slingo, J. M., 1987: The development and verification of a cloud prediction scheme for the ECMWF model. *Quart. J. Roy. Meteor. Soc.*, **113**, 899-927.
- Stull, R. B., 1988: An introduction to boundary layer meteorology. Kluwer Academic Publishers, The Netherlands, 666pp.
- Tiedtke, M., 1983: The sensitivity of the time-mean large-scale flow to cumulus convection in the ECMWF model. *Proceedings of the ECMWF Workshop on Convection in Large-Scale Models*, Reading, United Kingdom, ECMWF, 297-316.
- Tiedtke, M., 1989: A comprehensive mass flux scheme for cumulus parameterization in large-scale models. *Mon. Wea. Rev.*, **117**, 1779-1800.
- Yamada, T., and G. Mellor, 1975: A simulation of the Wangara atmospheric boundary layer data. *J. Atmos. Sci.*, **32**, 2309-2329.

Thanks for your attention !

Modeling is to understand what is
happening in nature !