

WRF Data, Utilities & Post-processing

Kelly Werner

July 2019



This material is based upon work supported by the National Center for Atmospheric Research, which is a major facility sponsored by the National Science Foundation under Cooperative Agreement No. 1852977.

Input Data



Input Data: Mandatory Fields

■ 3D Data (for data on constant pressure levels)

Temperature
U and V components of wind
Geopotential height
Relative Humidity (or Specific Humidity)

■ 3D Data (for data on native model levels)

Temperature
U and V components of wind
Geopotential height
Relative Humidity (or Specific Humidity)
Pressure

■ 2D Data

Surface pressure
Mean sea-level pressure
Skin temperature/SST
2 meter temperature and relative humidity
10 meter U and V components of wind
Soil data (temperature and moisture) and soil height

* Recommended Fields

LANDSEA mask field for input data
Water equivalent snow depth
SEAICE
Additional SST data



External Data Sources: Global

Name	Resolution	Coverage	Temporal Availability	Website
NCEP/NCAR Reanalysis (R1/NNRP)	209 km 6-hourly	Global	Jan 1948 - present	http://rda.ucar.edu/datasets/ds090.0
NCEP/DOE Reanalysis (R2)	209 km 6-hourly	Global	Jan 1979 - present	http://rda.ucar.edu/datasets/ds091.0
ERA Interim Data	1.125° - 0.703° 6-hourly	Global	Jan 1979 - present	http://rda.ucar.edu/datasets/ds627.0
ECMWF's Operational Model Analysis	Varying		Jan 2011 - present	http://rda.ucar.edu/datasets/ds113.0
NCEP GDAS/FNL Reanalysis	0.25° 6-hourly	Global	July 2015 - present	http://rda.ucar.edu/datasets/ds083.3
GFS Real-time	1°	Global		ftp://ftpprd.ncep.noaa.gov/pub/data/nccf/com/gfs
NCEP GFS/FNL Reanalysis	1° 6-hourly	Global	Aug 1999 - present	http://rda.ucar.edu/datasets/ds083.2
GFS Gridded Model Data	0.5° 24-hourly	Global	Dec 2002 - present	http://rda.ucar.edu/datasets/ds335.0
NCEP GFS 0.25°	0.25° 3-hourly & 12-hourly	Global	Jan 2015 - present	http://rda.ucar.edu/datasets/ds084.1

External Data Sources: North America

Name	Resolution	Coverage	Temporal Availability	Website
NAM Real-time	32/12 km 6-hourly	North America		ftp://ftpprd.ncep.noaa.gov/pub/data/nccf/com/nam
NAM Analysis	12 km 6-hourly	North America	Jan 2012 - present	http://rda.ucar.edu/datasets/ds609.0
GCIP NCEP Eta	40 km 3-hourly & 6-hourly	North America	April 1995 - present	http://rda.ucar.edu/datasets/ds609.2
NCEP NARR	32 km 3-hourly	North America	Nov 1979 - present	http://rda.ucar.edu/datasets/ds608.0



External Data Sources: Climate

Name	Resolution	Coverage	Temporal Availability	Website
NCEP Climate Forecast System Reanalysis (CFRS)	0.3° to 2.5° 6-hourly	Global	Jan 1979 - Dec 2010	http://rda.ucar.edu/datasets/ds093.0
NCEP Climate Forecast System Reanalysis II (CFRSv2)	0.2° to 2.5° 6-hourly	Global	Jan 2011 - present	http://rda.ucar.edu/datasets/ds094.0
NCAR CESM CMIP5 data (netCDF format)	6-hourly	Global	Jan 1950 - 2100	http://rda.ucar.edu/datasets/ds316.0
NCAR CESM CMIP5 data (IM - Bias Corrected)	6-hourly	Global	Jan 1951 - 2100	http://rda.ucar.edu/datasets/ds316.1
SST DATA				
NCEP SST Analysis	1° - 1/12°	Global		http://polar.ncep.noaa.gov/sst
NOMAD3 SST	1° - 0.25°	Global	Jan 1854 - present (depending which product)	http://nomads.ncdc.noaa.gov/data.php
NCEP & NCDC Reconstructed SST	1° - 2°	Global	Jan 1854 - Dec 2015	http://rda.ucar.edu/datasets/ds277.0



External Data Sources: RDA

<http://rda.ucar.edu>




External Data Sources: RDA

http://www2.mmm.ucar.edu/wrf/users/download/free_data.html



External Data Sources: NOMADS

<http://nomads.ncdc.noaa.gov>



NAM
GFS
RUC
CFS
NARR
R1/R2
SST

Utilities

- Grib and Intermediate Data
- Designing a model domain
- netCDF tools
- Other Utilities
- ImageMagick
- Special WRF Output Variables
- OBSGRID
- MET

GRIB Data Handling

▪ Documents

- <https://rda.ucar.edu/index.html#gribdoc> (GRIB1 data)
- <https://rda.ucar.edu/index.html#grib2doc> (GRIB2 data)

▪ Decoders

- *wgrib*, *wgrib2*, *unpackgrib2.c*, *grib2to1.c*
<http://rda.ucar.edu/#!/GRIB>
<http://www.cpc.ncep.noaa.gov/products/wesley/wgrib.html>
<http://www.cpc.ncep.noaa.gov/products/wesley/wgrib2>
- *g1print.exe* and *g2print.exe*
 - Show data available in GRIB1 and GRIB2 files
 - Available from util/ directory in WPS

GRIB Data Handling (cont'd)

▪ *grib2ctl.pl*

- Create .ctl and .idx files, so that you can plot GRIB files with GrADS
- <http://www.cpc.ncep.noaa.gov/products/wesley/grib2ctl.html>

▪ *ncl_convert2nc*

- Converts from grib format to netcdf format
http://www.ncl.ucar.edu/Document/Tools/ncl_convert2nc.shtml

Writing Intermediate File Format

- http://www2.ucar.edu/wrf/users/docs/user_guide/V4/v4.0/users_guide_chap3.html#Writing_Meteorological_Data

▪ wrf_wps_write_int

```
FIELD = "SST"
UNITS = "K"
DESC = "Sea Surface Temperature"
```

```
opt = True
opt@map_source           = "ERA-I Data"
opt@projection            = 0
opt@startloc              = "SWCORNER"
opt@startlon              = 0.0
opt@startlat              = -90.0
opt@deltalon              = 1.25
opt@deltalat              = 0.942408
opt@is_wind_earth_relative = False
opt@date                  = "2015-07-26_00:00:00"
opt@level                  = 200100.
```

```
wrf_wps_wrtie_int(IM_name,FIELD,UNITS,DESC,VAR(:,,:),opt)
```

Fortran script to convert netCDF to Intermediate format:

http://www2.mmm.ucar.edu/wrf/users/special_code.html



Reading Intermediate Format Files

NCL Code

▪ wrf_wps_read_int

! opens file

```
istatus = wrf_wps_open_int(filename)
```

! reads header

```
wrf_wps_rhead_int(istatus,head_real,field,
hdate, \
units,map_source,desc)
```

! reads slab

```
Slab = wrf_wps_rddata_int(istatus,nx,ny)
```

! Loop until reaching the end of the file

Found in WPS/util/

▪ rd_intermediate


```
=====
FIELD = TT
UNITS = K DESCRIPTION = TEMPERATURE
DATE = 2000-01-24_12:00:00 FCST = 0.000000
SOURCE = unknown model from NCEP GRID 212
LEVEL = 200100.000000
I,J DIMS = 185, 129
IPROJ = 1
REF_X, REF_Y = 1.000000, 1.000000
REF_LAT, REF_LON = 12.190000, -133.459000
DX, DY = 40.635250, 40.635250
TRUELAT1 = 25.000002
DATA(1,1) = 295.910950
=====
```



Utility: plotfmt

- The plotfmt program plots the fields in the ungribbed intermediate files

```
ncl plotfmt.ncl 'filename="FNL:2007-09-15_00"'
```



Found in WPS/util/




Plot Intermediate Files in netCDF Format

- Use the utility **int2nc.exe**
 - Converts intermediate files created by ungrib.exe to netcdf format
 - `./int2nc.exe FILE:yyyy-mm-dd_hh`


- To plot: **plotfmt_nc.ncl**

```
ncl plotfmt_nc.ncl 'inputFILE="FNL:YYYY-MM-DD_HH.nc"'
```

Plot Using ncview



Plot Using plotfmt_nc.ncl



*Both are found in WPS/util/



Model Domain Design

```
mpres@mpFillColor =
(/"background","DeepSkyBlue",
"ForestGreen","DeepSkyBlue",
"transparent"/)

mpres@mpGridSpacingF = 45


lnres@domLineColors = (/
"white", "Red", "Red", "Blue" /)

mpres@mpOutlineBoundarySets
"NoBoundaries" ; "Geophysical"
"National" ; "USStates"
"GeophysicalAndUSStates"
"AllBoundaries"

pares = True
pmres@gsMarkerColor = "White"
pmres@gsMarkerIndex = 16
pmres@gsMarkerSizeF = 0.01
gsn_polymarker(wks,mp,-
77.26,38.56,
pmres)
```

- **plotgrids.ncl**
 - WPS/util/plotgrids.ncl
 - Reads namelist information to generate plot
 - X11, png, pdf

Test Domain




Model Domain Design

```
DOMS = 1
DX = 36.
MAP = "mercator"
LAT1 = (/ -35.0, -45., -27. /)
LAT2 = (/ 0., -20., -23. /)
LON1 = (/ 131., 121., 125. /)
LON2 = (/ 171., 159., 131. /)
parent_id = (/ 0, 1, 2 /)
parent_grid_ratio = (/ 1, 3, 3 /)
```

Suggested namelist options

```
parent_id = 0,
parent_grid_ratio = 1,
i_parent_start = 1,
j_parent_start = 1,
e_we = 123,
e_sn = 107,
dx = 36000,
dy = 36000,
map_proj = 'mercator',
ref_lat = -17.50,
ref_lon = 151.00,
truelat1 = -17.00,
truelat2 = 0.00,
stand_lon = 151.00,
```



design_grids.ncl

http://www2.mmm.ucar.edu/wrf/users/special_code.html

netCDF Tools

NC0 Tools

<http://nco.sourceforge.net>

- netCDF Operators are command-line programs that take netCDF (HDF and/or DAP) files as input, then operate (e.g., derive new data, compute stats, print, manipulate metadata) and output to the screen or files in various formats (text, binary, netCDF, etc.)
- **ncdiff**
 - Shows the differences between 2 files
 - `ncdiff input1.nc input2.nc diff.nc`
- **ncrcat (nc cat)**
 - Writes specified variables/times to a new file, or concatenates files
 - `ncrcat file1.nc file2.nc combined.nc`
 - `ncrcat -d Time,0,231 -v RAINNC wrfout* RAINNC.nc`
- **ncra (nc average)**
 - Averages variables in files and writes to a new file
 - `ncra -v T2 file1.nc file2.nc -o T2.nc`
 - `ncra -v T2 wrfout* -o T2.nc`

NCO Tools (continued)

<http://nco.sourceforge.net>

■ **ncrename**

- Renames variables, dimensions, attributes

```
ncrename -v LANDUSE, LAND -a missing_value, _FillValue
file.nc
```

■ **ncks** (nc kitchen sink)

- Combination of several NCO tools to allow cutting/pasting subsets of data into a new file

- Extracting a specific variable

```
ncks -v RAINNC wrfout_d01_2015-06-01_00:00:00 RAINNC.nc
```

- Splitting files

```
ncks -d Time,1,1 wrfout -o wrfout1.nc
```




NCO Tools: Other Available Operators

- **ncap2**: arithmetic processor
- **ncatted**: ATtribute editor
- **ncbo**: binary operator (includes ncadd, ncsubtract, ncmultiply, ncdivide)
- **ncea**: ensemble averager
- **ncecat**: ensemble conCATenator
- **ncflint**: FiLe INterpolator
- **ncpdq**: permute dimensions quickly, pack data quietly
- **ncwa**: weighted averager



ncview



http://meteora.ucsd.edu/~pierce/ncview_home_page.html



- A graphical interface that allow quick viewing of netCDF files
 - All variables found in file
 - Detect where things go wrong
- Other options
 - Time series
 - Vertical Cross Section
- WRF/WPS files
 - Any netCDF format file
 - geo_em.d0*, met_em.d0*, wrfinput_d0*, wrfout.d0*, wrfrst.d0*



ncview




- Beginning WRFV3.7
- Works with wrfinput* and wrfout* files
- Must have 1 time period per file



ncBrowse

<http://www.epic.noaa.gov/java/ncBrowse/>



ncdump

- Reads a netCDF dataset and prints information from that dataset
- **ncdump -h file**
 - Prints header (inclusive list of variables in the file)
- **ncdump -v VAR file**
 - Prints specific data for the variable 'VAR'
- **ncdump -v Times file**
 - Prints the times that are included in the file



ncdump -v Times file

```
netcdf wrfout_d01_2000-01-24_12:00:00 {
dimensions:
    Time = UNLIMITED ; // (3 currently)
    DateStrLen = 19 ;
    west_east = 73 ;
    south_north = 60 ;
    west_east_stag = 74 ;
    bottom_top = 27 ;
    south_north_stag = 61 ;
    bottom_top_stag = 28 ;
variables:
    char Times(Time, DateStrLen) ;
    float LU_INDEX(Time, south_north, west_east) ;
        LU_INDEX:FieldType = 104 ;
        LU_INDEX:MemoryOrder = "XY" ;
        LU_INDEX:description = "LAND USE CATEGORY" ;
        LU_INDEX:units = "" ;
        LU_INDEX:stagger = "" ;
.....
global attributes:
    :TITLE = " OUTPUT FROM WRF V3.4.1 MODEL";
    :START DATE = "2000-01-24 12:00:00" ;
    :WEST-EAST GRID DIMENSION = 74 ;
    :SOUTH-NORTH GRID DIMENSION = 61 ;
    :BOTTOM-TOP GRID DIMENSION = 28 ;
    :DX = 30000.f ;
    :DY = 30000.f ;
.....
data:
    Times =
        "2000-01-24 12:00:00",
        "2000-01-24 18:00:00",
        "2000-01-25 00:00:00"
```



Other Utilities

- Additional utilities
 - **read_wrf_nc**: reads WRF netCDF file, outputs various data
 - **iowrf**: extracts a box from WRF netCDF files, thin or destagger data
 - **wrf_interp**: interpolates WRF output files to pressure, height-agl, height-msl, potential temp, and equivalent potential temp, and can perform underground extrapolation
 - **p_interp**: converts wrfout data (from sigma levels) to pressure levels
 - **v_interp**: adds vertical levels in WRF input and boundary files
 - **diffwrf**: performs several functions, including making comparisons of two WRF files (available in WRF/external/io_netcdf/)
 - For more details on the above utilities, see:
<http://www2.mmm.ucar.edu/wrf/users/utilities/util.htm>
- To download utilities:
http://www2.mmm.ucar.edu/wrf/users/download/get_sources.html



ImageMagick

<http://www.imagemagick.org>

- Converts graphical files from one format to another

```
convert file.pdf file.png
convert file.png file.bmp
```
- Many options available
 - Rotate frames, trim white space, etc.
 - 2 ways to use
 - 1) `display plot.png`
 - 2) `Convert -trim +repage -background white -flatten plot.pdf plot.png`
- Can make movies
 - Can create individual frames for each image
- Maintains high resolution - great for publishing!
- Cannot deal with .ncgm files



Special WRF Output Variables

- The WRF model outputs the state variables defined in the Registry file, and these state variables are used in the model's prognostic equations. Some of these variables are perturbation fields and therefore, the following definitions for reconstructing meteorological variables are necessary:

Total geopotential	PH + PHB
Total geopotential height in m	(PH + PHB) / 9.81
Total potential temp in K	T + 300
Total pressure in mb	(P + PB) * 0.01
Wind components, grid relative	U, V
Surface pressure in Pa	Psfc
Surface winds, grid relative	U10, V10 (valid at mass points)
Surface temp and mixing ratio	T2, Q2

See [WRF/Registry/Registry.EM_COMMON](#) for description of variables



OBSGRID




OBSGRID

- To improve a first-guess gridded analysis by incorporating additional observational information
 - Traditionally first-guess analysis came from low-resolution global analysis and forecast grids
 - These days, higher-resolution, regional scale analyses are more readily available
- When is this method useful?
 - When using very coarse resolution first-guess input data
 - If you conducted a field campaign and have acquired very high-resolution station data (for example)



OBSGRID: Basic Concept



OBSGRID: How to Run

- Get the source code
<http://www2.mmm.ucar.edu/wrf/users/downloads.html>
- Compile (serially)
- Prepare observation files
- Edit the namelist.oa
- Link in met_em* files from WPS
- Run the program
 - ./obsgrid.exe
- Check your output

See the WRF Users' Guide for detailed information:

http://www2.mmm.ucar.edu/wrf/users/docs/user_guide_V4/v4.0/users_guide_chap7.htm



OBSGRID: Output

- **metoa_em***
 - Final analysis files at sfc and pressure levels
 - Can be used to replace met_em* files during real.exe
- **wrfsfdda_d0***
 - Contains sfc analyses for T, TH, U, V, RH, QV, PSFC, PMSL, and a count of obs within 250 km of each grid point
 - Use as input for surface analysis nudging (in &fdda section of namelist, set grid_fdda = 1 and grid_sfdda = 1)
- **OBS_DOMAIN*01** (e.g., OBS_DOMAIN101, OBS_DOMAIN103)
 - Contain list of all observations available to OBSGRID program
 - User has option to set values > a specified value to "missing"
 - Used during WRF for observational nudging
 - Must all be concatenated into a single file (OBS_DOMAIN101)
 - Use 'run_cat_obs_files.csh' to do this



OBSGRID: Output (continued)

- **qc_obs_raw.d01.YYYY-MM-DD_HH:mm:ss.tttt(.nc)**
 - Similar to OBS_DOMAINXXX, but raw data, and output is in either ASCII or netCDF format
 - ASCII file can be used as input to plotting utility "plot_sounding.exe"
 - NetCDF file can be used to plot both station & sounding data
- **qc_obs_used.d01.YYYY-MM-DD_HH:mm:ss.tttt(.nc)**
 - Same as above (can be used in the same way), but identical data to OBS_DOMAIN*01 files
- **qc_obs_used_earth_relative.d01.YYYY-MM-DD_HH:mm:ss.tttt(.nc)**
 - Identical to above, but winds are in an earth-relative framework, rather than model-relative
 - The non-netCDF version can be used as input to MET verification software
- **plotobs_out.d01.YYYY-MM-DD_HH:mm:ss.tttt**
 - Lists data by variable and by level
 - Each observation used for objective analysis is grouped with all associated observations for plotting or other diagnostic purposes



OBSGRID: Plot Utilities

- **sounding.ncl (found in util/)**
 - Generates & plots soundings from qc_obs_raw* and qc_obs_used*
- **station.ncl (found in util/)**
 - creates station plots for each analysis level



MET Verification Software

- Model Evaluation Tools (MET)
- Provides all the basics (e.g., RMSE, bias, skill scores)
- Provides
 - Advanced spatial methods (wavelets, objects)
 - Confidence intervals
- Download it
<http://www.dtcenter.org/met/users/downloads/>
- Support
met_help@ucar.edu
- Documentation
<http://www.dtcenter.org/met/users/docs/overview.php>



Post-processing

- Supported Packages
- ARWpost
- RIP4



Supported Post-processing Packages

http://www2.mmm.ucar.edu/wrf/users/docs/user_guide_V3/contents.html

Package	Users' Guide Page #	Information
NCL	9-2	Graphical package supported by NCAR/CISL (http://forum.mmm.ucar.edu/ and ncl-talk@ucar.edu)
ARWpost	9-29	Converter (GrADS); ARWpost supported by wrfhelp: http://forum.mmm.ucar.edu/
RIP4	9-20	Converter and interface to graphical package, NCAR graphics, supported by wrfhelp: http://forum.mmm.ucar.edu/
UPP	9-36	Converter (GrADS & GEMPAK) (upp-help@ucar.edu)
VAPOR	9-38	Converter and graphical package Supported by VAPOR (vapor@ucar.edu)
IDV	None - see unidata.ucar.edu	GRIB (from UPP) GEMPAK (from wrf2gem) Vis5d CF compliant data (from wrf_to_cf) Supported by unidata (support@unidata.ucar.edu)
GEMPAK	None - see: unidata.ucar.edu/software/gempak	Data from wrf2gem or UPP Supported by unidata (support@unidata.ucar.edu)



Choosing the Right Tool

- Can it read your data?
- Will you need to pre-process the data first?
- Is it purely a visualization tool, or does it include post-processing?
- Can it handle big datasets?
- Which diagnostic/statistical functions does it have?
- How easy is it to add diagnostics?
- 3D or 2D visualization?
- Can it handle staggered grids?
- How is data below the ground handled?
- Vertical grids?
- How are model time stamps handled?
- Easy to use?
- Cost of package?
- How well supported is it?



Data Handling

	NCL	RIP4	GrADS	UPP	VAPOR	IDV
netCDF		ripdp	ARWpost	converter	converter	converter
GRIB						
ASCII						
shapefiles						
geogrid & metgrid output						
intermediate file format	V6.2.0 V6.3.0					
wrfinput data						
Idealized data						
wrfoutput						
big data						




Post-processing

	NCL	RIP4	GrADS	UPP	VAPOR	IDV
Post-processing						
Data output						
3D						
diagnostics	some	a lot	some	some	limited	limited
Add diagnostics	Very easy	easy	easy	Relatively easy	Not as easy	Not as easy
Vertical output Coordinate	Model pressure height	Model pressure height	Model pressure height	pressure	model	model
Extrapolate Below ground						



Model Staggering

Why is a converter necessary if a package can display netCDF files?



- 1) Converter co-locates data to mass points
- 2) Converter translates variables - e.g., "T" is not really temp. Must add 300 for actual temp (K)



ARWpost




ARWpost: General Information


- Converter
 - Reads in wrf-arw model data, creates GrADS output files
 - Requires GrADS to display
- Download Code
 - http://www2.mmm.ucar.edu/wrf/users/download/get_sources.html
- GrADS software is only needed to display data, not needed to compile the code
 - <http://www.iges.org/grads/grads.html>
- Online Tutorial
 - <http://www2.mmm.ucar.edu/wrf/OnLineTutorial/Graphics/ARWpost/index.php>
- Generate a number of graphical plots
 - Horizontal
 - Cross-section
 - skewT
 - Meteogram
 - Panel



ARWpost: Example Plots



ARWpost: Example Functions



ARWpost: Diagnostics

- cape - 3d cape
- cin - 3d cin
- mcape - maximum cape
- mcin - minimum cin
- clfr - low/middle/high cloud fraction
- dbz - 3d reflectivity
- max_dbz - maximum reflectivity
- geopt - geopotential
- height - model height in km
- lcl - lifting condensation level
- lfc - level of free convection
- pressure - full model pressure in hPa
- rh - relative humidity
- rh2 - 2 m relative humidity
- theta - potential temperature
- tc - temperature in degrees C
- tk - temperature in degrees K
- td - dew point temperature in degrees C
- td2 - 2m dew point temperature in degrees C
- slp - sea level pressure
- umet & vmet - winds rotated to Earth coordinates
- u10m & v10m - 10 m winds rotated to Earth coordinates
- wdir - wind direction
- wspd - wind speed coordinates
- wd10 - 10 m wind direction
- ws10 - 10 m wind speed



ARWpost: Scripts

Script Name	Description
cbar.gs	Plots a color bar on shaded plots
rgbset.gs	Allows you to add/change colors from color # 20 - 99
skew.gs	Program to plot a skewT
plot_all.gs	Automatically finds all .ctl files in the directory and lists them so the user can pick when to use, will plot all fields chosen
rain.gs (real data only)	Plots total rainfall (must have data that contain fields RAINC and RAINNC)
cross_z.gs (real data only)	Plots a NS and EW cross section of RH and T (C)



RIP4




RIP4: General Information


- Requires NCAR Graphics Libraries
 - <http://www.ncl.ucar.edu>
- Source Code
 - http://www2.mmm.ucar.edu/wrf/users/download/get_source.html
- Documentation
 - Included in program's tar file (in Doc/ directory)
 - <http://www2.mmm.ucar.edu/wrf/users/docs/ripug.htm>
- Online Tutorial
 - <http://www2.mmm.ucar.edu/wrf/users/graphics/RIP4/RIP4.htm>




RIP4: Example Plots



RIP4: Example Plots



RIP4: Program Flow



RIP4: Namelist (&userin)

- Use namelist to control
 - processing times, intervals, title information, text quality on a plot
 - whether to do time series, trajectory, or to write output for Vis5D
 - Full explanation for namelist variables is available in the user document
- ptimes, ptimeunits - times to process
- tacc - tolerance for processing data
- iusedaylightrule - 1 applied, 0 not applied
- idotser - generate time series output
- icgmsplit - split metacode into several files
- itrajcalc - 0, 1 ONLY when doing trajectory calculations
- rip_root - override RIP_ROOT
- ncarg_root - output type: X11, cgm, pdf, ps

RIP4: Common Error Message

GKS ERROR NUMBER 2 ISSUED FROM SUBROUTINE
GCLKS :--GKS NOT IN PROPER STATE: GKS SHALL BE IN
STATE GKOPFORTRAN STOP

- Usually NOT a graphics error.
- More often this is an error with the times you are asking RIP to process
 - Check the ptimes in your .in file
 - Check the xtimes files created by RIPDP



Questions?

