

# *Considerations for Designing an Numerical Experiment*

*Wei Wang*


*NCAS/NCAR Tutorial, Lincoln UK*

*October 2019*



# Issues

- Domain configuration
- Horizontal and vertical grid sizes
- Input data (both static and meteorological)
- Model options





# Domains

- In general,
  - IC is more important for simulations of a few days;
  - BC is more important for longer simulations.
- How large do they need to be?
  - Should not be too small, otherwise solution will be determined by forcing data
  - No less than 100x100 (at least 10 grid points are in the boundary zone)
- Where to place my lateral boundaries?
  - Avoid steep topography
  - Away from the area of interest




# *Note on Configuring Domains: Horizontal*





# *Note on Configuring Domains:*

## *Effect of domain sizes*

Large regional domain




Smaller regional domain



*(From Warner, 2011)*


# Note on Configuring Domains: Effect of lateral boundary conditions



(From Gaudet et al. WRF Users' Workshop 2012, talk 3.5)


# Domains

- How many vertical levels should I use?
  - At least 30 or more levels for model top at 50 mb
  - 50 mb or higher model top is recommended
  - Vertical grid distance should not be larger than 1000 m:
    - Radiation, microphysics, less accurate lateral BC
  - Related to horizontal grid size too: if finer horizontal grid size is used, consider adding a few more levels in the vertical



# Note on Configuring Domains:

## Vertical levels



# Domains

- Consider the placement of your domains:
  - What map projection to use?
  - Check the range of the map scale factor after running *geogrid*
    - Values should be close to 1

\* Placement of the domain will affect the time step used in the model.




# Nests:

- When should I use nests?

Some of the reasons may be:

- Input data resolution is too coarse
- Input data may not be adequate as LBC
- There isn't sufficient computing resources


- Nest domain sizes should not be too small;
- Nest boundary should be kept away from coarse domain boundary, and steep topography;
- If you use a nest, do not save on coarse domain – it's cheap (and may scale better when using large number of processors)



# Input Data

- Check static data:  
e.g. landuse: *does it represent my area well?*
- Know about the data: *how good are the data?*
  - Forecast data
  - Reanalysis data
  - Climate model data
- How frequent do I need to have boundary conditions?
  - More frequent is better

\* Good data will go a long way to ensure good outcome.



# *Model Options*

---

- What do I start with?
  - What other people have success with?
    - References, papers
    - Consider well-tested options first
  - Simple options first:

For example,

    - Graupel may not be important if  $dx \gg 10$  km
    - mixed layer ocean model may not be needed if the modeled track isn't correct
    - Use analyses from weather centers before trying to create your own (via either *obsgrid* or data assimilation) for both initial and lateral boundary conditions
    - Single domain first, before using many nests



# *Model Options*

---

- Choose physics for appropriate grid sizes
  - Use a cumulus scheme if grid size  $> 10$  km
  - A cumulus scheme isn't needed when grid size  $< 4$  km
  - Avoid grid sizes 5 – 10 km if not using scale-aware CPS
  - Use a PBL for grid size  $> 500$  m
  - Use LES options for grid size  $< 100$  m
- Consider other options:


For example,

  - Upper level damping (damp\_opt = 3)
  - Gravity-wave drag (gwd\_opt = 1)
  - Slope effect on radiation when grid size  $< 2$  km (slope\_rad = 1)



# Verification:

- Important to verify:
  - Knowing where model is biased can be very useful
- Verifying high-resolution model can be tricky:
  - e.g. phase error, which punishes higher resolution model more
  - Neighborhood method more appropriate




# A forecast example

-- What can we learn from this example?




# Derecho forecast from NCAR's 2012 RT

Observed  
radar  
composite




(from NOAA/SPC)

Forecast max-  
column  
reflectivity from  
3 km model,  
starting from  
1200 UTC  
June 29



IC: Fully  
cycled  
analysis  
starting from  
late April  
using WRF-  
DART



IC

F06

F09




F12

1800 UTC, 2 pm EDT




2100 UTC, 5 pm EDT

0000 UTC, 8 pm EDT





EnKF



NAM




GFS




# *Sensitivity to physics and initial conditions:*


29 June 2012  
Derecho




DART-Morrison-  
MYJ




DART-Morrison-YSU




DART-Thompson-MYJ




GFS



NAM



RUC




scal



(From Weisman)

# Resolution Differences: simulated max winds


3 km results



15 km results



# NCAR Real-time Forecast Domain (2013)





# What this case show:

- Initial conditions are important
- Different initial conditions will likely give different solutions
- Compared to model runs using different physics options, changing initial conditions is likely to have larger impact
- Model resolution matters



# Another example of IC impact




Five day forecast of hurricane Joaquin from 0000 UTC  
9/30/2016



# An example of nest feedback

00z SST, TSK  
Fcst: 60 h  
Total precip. in past 6 h


Init: 06 UTC Sun 04 May 14  
Valid: 18 UTC Tue 06 May 14 (02 LST Wed 07 May 14)



feedback = 0

e04: feedback=1. 00z SST, TSK  
Fcst: 60 h  
Total precip. in past 6 h

Init: 06 UTC Sun 04 May 14  
Valid: 18 UTC Tue 06 May 14 (02 LST Wed 07 May 14)




feedback = 1

60-h forecast of the 6-h rainfall for the period ending  
18 UTC 6 May 2014 (courtesy of J. Bresch)



# Bottomline..

- Model results can be affected by many choices:
  - Domain configuration, both horizontal and vertical;
  - Input data;
  - Initial and lateral boundary conditions.
- Model has limitations:
  - Physics: biases, may not represent certain process well, etc.
  - Limitation of the lateral boundaries
- **Always check the output after each program**



## ***Other Best Practice Reading:***

- *“12 steps toward improving the outcome” by C. Davis:*  
<http://www2.mmm.ucar.edu/wrf/users/workshops/WS2012/ppts/discussion1.pdf>
- *“WRF Advanced usage and Best Practices” by Dudhia and Wang:*  
[http://www2.mmm.ucar.edu/wrf/users/workshops/WS2014/ppts/best\\_prac\\_wrf.pdf](http://www2.mmm.ucar.edu/wrf/users/workshops/WS2014/ppts/best_prac_wrf.pdf)



## ***References:***

Numerical Weather and Climate Prediction, 2011. By Thomas Warner, *Cambridge University Press*.

Warner, T., 2011. Quality assurance in atmospheric modeling. *Bull. Amer. Met. Soc. Dec. issue, p1601 – 1611.*

Stensrud, D., 2007. Parameterization Schemes: Keys to Understanding Numerical Weather Prediction Models. *Cambridge University Press*.

Haltiner G. and R. Williams, 1980. Numerical Prediction and Dynamic Meteorology. *Wiley*.

